

1256904

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office

December 06, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/523,600
FILING DATE: *November 20, 2003*
RELATED PCT APPLICATION NUMBER: PCT/US04/37181

Certified by




Jon W Dudas

Acting Under Secretary of Commerce
for Intellectual Property
and Acting Director of the U.S.
Patent and Trademark Office

BEST AVAILABLE COPY

PROVISIONAL APPLICATION FOR PATENT COVER SHEET


This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c)

Docket Number		P-16541	Type a plus sign (+) inside this box ->	+	
INVENTOR(s)/APPLICANT(s)					
LAST NAME	FIRST NAME	MIDDLE NAME	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)		
WARSHAWSKY	ALAN	M	CARMEL, INDIANA		
LU	JIANLIANG		FISHERS, INDIANA		
NAGPAL	SUNIL		CARMEL, INDIANA		
MA	TIANWEI		CARMEL, INDIANA		
SHEN	QUANRONG		FISHERS, INDIANA		
TITLE OF THE INVENTION (280 characters max)					
Vitamin D Receptor Modulators					
CORRESPONDENCE ADDRESS					
Eli Lilly and Company Patent Division P.O. Box 6288 Indianapolis, Indiana 46206-6288			 25885 PATENT TRADEMARK OFFICE		
STATE	IN	ZIP CODE	46206-6288	COUNTRY	USA
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification	Number of pages	192	<input type="checkbox"/> Small Entity Statement		
<input type="checkbox"/> Drawing(s)	Number of Sheets		<input type="checkbox"/> Other (Specify)		
METHOD OF PAYMENT (check one)					
<input type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fees			PROVISIONAL FILING FEE AMOUNT (\$)	\$160.00	
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit Deposit Account Number:	05-0840				

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.
☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,
SIGNATURE


Roger S. Benjamin

Date 20/ 11 / 2003


REGISTRATION NO.
(if appropriate)

27,025

TYPED or PRINTED NAME

☐ Additional inventors are being named on separately numbered sheets attached hereto

PROVISIONAL APPLICATION FOR PATENT FILING ONLY

"Express Mail" mailing label number	ER 357 404840 US	Date of Deposit	20 November 2003
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to the Commissioner for Patents, P.O. Box 1450, Alexandria, VA, 22313-1450.			
Printed Name	QUEEN THOMAS	Signature	

VITAMIN D RECEPTOR MODULATORS**BACKGROUND OF THE INVENTION**

Vitamin D₃ Receptor (VDR) is a ligand dependent transcription factor that belongs to the superfamily of nuclear hormone receptors. The VDR protein is 427 amino acids, with a molecular weight of ~50 kDa. The VDR ligand, 1 α ,25-dihydroxyvitamin D₃ (the hormonally active form of Vitamin D) has its action mediated by its interaction with the nuclear receptor known as Vitamin D receptor ("VDR"). The VDR ligand, 1 α ,25-dihydroxyvitamin D₃ (1 α ,25(OH)₂D₃) acts upon a wide variety of tissues and cells both related to and unrelated to calcium and phosphate homeostasis.

The activity 1 α ,25-dihydroxyvitamin D₃ in various systems suggests wide clinical applications. However, use of conventional VDR ligands is hampered by their associated toxicity, namely hypercalcemia (elevated serum calcium). Currently, 1 α ,25(OH)₂D₃, marketed as Rocaltrol® pharmaceutical agent (product of Hoffmann-La Roche), is administered to kidney failure patients undergoing chronic kidney dialysis to treat hypocalcemia and the resultant metabolic bone disease. Other therapeutic agents, such as Calcipotriol® (synthetic analog of 1 α ,25(OH)₂D₃) show increased separation of binding affinity on VDR from hypercalcemic activity.

Chemical modifications of 1 α ,25(OH)₂D₃ have yielded analogs with attenuated calcium mobilization effects (R. Bouillon et. al., Endocrine Rev. 1995, 16, 200-257). One such analog, Dovonex ® pharmaceutical agent (product of Bristol-Meyers Squibb Co.), is currently used in Europe and the United States as a topical treatment for mild to moderate psoriasis (K. Kragballe et. al., Br. J. Dermatol. 1988, 119, 223-230).

Other Vitamin D₃ mimics have been described in the publication, Vitamin D Analogs: Mechanism of Action of Therapeutic Applications, by Nagpal, S.; Lu, J.; Boehm, M. F., Curr. Med. Chem. 2001, 8, 1661-1679.

"Express Mail" mailing label number EP 357404840 US
Date of Deposit: 20 November 2003

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 110 on the date indicated above and is addressed to the Commissioner for Patent, P.O. Box 1450, Alexandria, VA. 22313-1450

Queen Thomas

(Typed or Printed name of person mailing paper)

Queen Thomas

(Signature of person mailing paper or fee)

Although some degree of separation between the beneficial action and calcium raising (calcemic) effects has been achieved with these VDR ligands, to date the separation has been insufficient to allow for oral administration to treat conditions such as osteoporosis, cancers, leukemias, and severe psoriasis.

One example of a major class of disorder that could benefit from VDR mediated biological efficacy in the absence of hypercalcemia is osteoporosis. Osteoporosis is a systemic disorder characterized by decreased bone mass and microarchitectural deterioration of bone tissue leading to bone fragility and increased susceptibility to fractures of the hip, spine, and wrist (World Health Organization WHO 1994). Osteoporosis affects an estimated 75 million people in the United States, Europe, and Japan.

Within the past few years, several antiresorptive therapies have been introduced. These include bisphosphonates, hormone replacement therapy (HRT), a selective estrogen receptor modulator (SERM), and calcitonins. These treatments reduce bone resorption, bone formation, and increase bone density. However, none of these treatments increase true bone volume nor can they restore lost bone architecture.

Another major disorder that could benefit from VDR mediated biological activity is psoriasis. Psoriasis is one of the most common dermatologic diseases and is a chronic inflammatory skin condition characterized by erythematous, sharply demarcated papules and rounded plaques, covered by silvery micaceous scale.


Synthetic VDR ligands with reduced calcemic potential have been synthesized. For example, a class of bis-phenyl compounds stated to mimic 1α , 25-dihydroxyvitamin D_3 is described in US Patent No. 6,218,430 and the article; "Novel nonsecosteroidal vitamin D mimics exert VDR-modulating activities with less calcium mobilization than 1α , 25-Dihydroxyvitamin D_3 " by Marcus F. Boehm, et. al., Chemistry & Biology 1999, Vol 6, No. 5, pgs. 265-275.

Synthetic VDR ligands having an aryl-thiophene nucleus are described in United States provisional patent application SN 60/384151, filed 29 May 2002.


There remains a need for improved treatments using alternative or improved pharmaceutical agents that mimic 1α , 25-dihydroxyvitamin D_3 to stimulate bone formation, restore bone quality, and treat other diseases without the attendant disadvantage of hypercalcemia.

SUMMARY OF THE INVENTION


Novel compounds having a nucleus of Formula "(TP)" or "(PT)" have been found effective as Vitamin D Receptor modulators (VDRM):



or




Compounds of the invention with VDR modulating activities are represented by formulae (IA)



wherein the variables R, R', RP, RP₃, LP₁, LP₂, Z_P, RB, RB', RT₃, L_{TB} and Z_{TB} are as hereinafter defined.

Compounds of the invention with VDR modulating activities are also represented by formulae (IB)



wherein the variables R, R', RP, RP₃, L_{P1}, L_{P2}, Z_P, RB₇, RB, RB₄, RT₃, L_{BT} and Z_{BT} are as hereinafter defined.

In another aspect, the present invention is directed towards pharmaceutical compositions containing pharmaceutically effective amounts of compounds of formulae IA or IB or a pharmaceutically acceptable salt or a prodrug thereof, either singly or in combination, together with pharmaceutically acceptable carriers and/or auxiliary agents.

Another aspect of the invention is a pharmaceutical formulation for treatment or prevention of osteoporosis containing pharmaceutically effective amounts of the vitamin D receptor modulator compound of formulae IA or IB alone or together with pharmaceutically effective amounts of co-agents conventionally used for the treatment of osteoporosis.

Another aspect of the invention is a pharmaceutical formulation for treatment or prevention of psoriasis containing pharmaceutically effective amounts of the vitamin D receptor modulator compound of formulae IA or IB alone or together with pharmaceutically effective amounts of co-agents conventionally used for the treatment of psoriasis.

Another aspect of the invention is a pharmaceutical formulation for treatment or prevention of prostate cancer containing pharmaceutically effective amounts of the vitamin D receptor modulator compound of formulae IA or IB alone or together with pharmaceutically effective amounts of co-agents conventionally used for the treatment of prostate cancer.

Another aspect of the invention is to use the compounds of formulae IA or IB to treat disease states responsive to Vitamin D receptor ligands.

Another aspect of the invention is the prevention and treatment of acne, actinic keratosis, alopecia, Alzheimer's disease, autoimmune induced diabetes, bone

fracture healing, breast cancer, Crohn's disease, prostate cancer, colon cancer, Type I diabetes, host-graft rejection, hypercalcemia, Type II diabetes, leukemia, multiple sclerosis, insufficient sebum secretion, osteomalacia, osteoporosis, insufficient dermal firmness, insufficient dermal hydration, myelodysplastic syndrome, psoriatic arthritis, psoriasis, renal osteodystrophy, rheumatoid arthritis, scleroderma, seborrheic dermatitis, skin cancer, systemic lupus erythematosus, skin cell damage from Mustard vesicants, ulcerative colitis, and wrinkles, by administering to a mammal in need thereof a pharmaceutically effective amount of a compound of Formulae IA or IB.

DETAILED DESCRIPTION OF THE INVENTION

Definitions:

The term, "abscess" refers to adverse complications often associated with surgery, trauma, or diseases that predispose the host to abscess formation from encapsulated bacteria lymphocytes, macrophages, and etc.

The term, "adhesion" refers to the adverse and abnormal union of surfaces normally separate by the formation of new fibrous tissue resulting from an inflammatory process.










The term, "compound of the invention" refers to a compound represented by Formulae IA or IB or as set out as products of the Examples or synthesis schemes described herein.

The term, "Active Ingredient" means a compound of the invention.

The term, "Mustard" is inclusive of both sulfur mustards and nitrogen mustards, either alone or in any combination. Exemplary of such compounds are the vesicants; bis(2-chloroethyl) sulfide (Chemical Agent Symbol HD), $\text{Cl}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{Cl}$ 1,2-bis(2-chloroethylthio)ethane (Chemical Agent Symbol Q), $\text{Cl}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{Cl}$; bis(2-chloroethylthioethyl) ether, $\text{Cl}(\text{CH}_2)_2\text{S}(\text{CH}_2)\text{O}(\text{CH}_2)_2\text{S}(\text{CH}_2)_2\text{Cl}$ (Chemical Agent Symbol T); tris(2-chloroethyl) amine (Chemical Agent Symbol HN3) $\text{N}(\text{CH}_2\text{CH}_2\text{Cl})_3$; N-methyl-2,2'-dichlorodiethylamine (Chemical Agent Symbol NH2); and 2,2'-dichlorotriethylamine, $\text{CH}_3\text{CH}_2\text{N}(\text{CH}_2\text{CH}_2\text{Cl})_2$ (Chemical Agent Symbol NH1).













The term, "(Acidic Group)" means an organic group that acts as a proton donor capable of hydrogen bonding. Illustrative of an (Acidic Group) is a group selected from the following:

-C(O)OH,
-5-tetrazolyl,




P-16541


7




or




The term, “-1,3-thiazolidine-2,4-dione-5-ethylidene”, refers to the radical represented by the structural formula:




The term, “-CH₂-C(O)-N-pyrrolidine” refers to the radical represented by the structural formula:




The term, “-CH₂-N-pyrrolidin-2-one” refers to the radical represented by the structural formula:




The term, “-CH₂-(1-methylpyrrolidin-2-one-3-yl)” refers to the organic radical represented by the structural formula:




The term, “1,3,4-oxadiazolin-2-one-5-yl” refers to the organic radical represented by the structural formula:




The term, “1,3,4-oxadiazolin-2-thione-5-yl” refers to the organic radical represented by the structural formula:



The term, "imidazolidine-2,4-dione-5-yl" refers to the organic radical represented by the structural formula:



The term, "isoxazol-3-ol-5-yl" refers to the organic radical represented by the structural formula:



The dotted line symbol crossing a solid line representing a bond




means that the bond so marked is the bond of attachment.


The term, "mammal" includes humans.

The term "halo" refer to fluorine, chlorine, bromine, and iodine.

The term "sulfonate" refers to the group




where R''' is C₁-C₅ alkyl, C₁-C₅ fluoroalkyl ,




where R' is -CO₂H, -CO₂R''', -OH, -CF₃, or C₁-C₅ alkyl.


The term "sulfonamide" refers to the group methyl, ethyl, branched C₃-C₅ alkyl,




where R'' is H, C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, or



or



where R''' is C₁-C₅ alkyl, C₁-C₅ fluoroalkyl,



and

where R' is -CO₂H, -CO₂R''', -OH, -CF₃, or C₁-C₅ alkyl.

The term, "C₁₋₃ alkyl" refers to an alkyl group selected from methyl, ethyl, n-propyl, and isopropyl.

The term, "branched C₃-C₅ alkyl" is an alkyl group selected from 1-methylethyl; 1-methylpropyl; 2-methylpropyl; 1,1-dimethylethyl; 1,1-dimethylpropyl; 1,2-dimethylpropyl; or 2,2-dimethylpropyl. Preferred branched C₃-C₅ alkyl groups are 2-methylpropyl and 1,1-dimethylethyl, with the 1,1-dimethylethyl group being most preferred.

The term "alkenyl" refers to aliphatic groups wherein the point of attachment is a carbon-carbon double bond, for example vinyl, 1-propenyl, and 1-cyclohexenyl. Alkenyl groups may be straight-chain, branched-chain, cyclic, or combinations

thereof, and may be optionally substituted. Suitable alkenyl groups have from 2 to about 20 carbon atoms.

The term "C₁-C₅ alkyl" refers to saturated aliphatic groups including straight-chain, branched-chain, and cyclic groups and any combinations thereof. Examples of C₁-C₅ alkyl groups are methyl, ethyl, n-propyl, from 1-methylethyl; n-butyl, 1-methylpropyl; 2-methylpropyl; 1,1-dimethylethyl; n-amyl, 1,1-dimethylpropyl; 1,2-dimethylpropyl; and 2,2-dimethylpropyl.

The term "cycloalkyl" includes organic radicals such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The term, "cycloalkenyl" includes organic radicals such as cyclopropenyl, cyclobutenyl, cyclopentenyl and cyclohexenyl.

The term, "C₁-C₅ fluoroalkyl" is an alkyl group containing fluorine and includes organic radicals such as -CF₃, -CHF₂, -CH₂F, -CF₂CF₃, -CHF₂CF₃, -CH₂CF₃,

-CH₂CHF₂, and -CH₂CH₂F, with -CF₃ being preferred.

The abbreviation, "Me" means methyl.

The abbreviation, "Et" means ethyl.

The abbreviation, "iPr" means 1-methylethyl.


The abbreviation, "tBu" means 1,1-dimethylethyl.

The term, "terminal hydroxyalkyl" is a group selected from


3-methyl-3-hydroxypentyl,
3-methyl-3-hydroxypentenyl,
3-methyl-3-hydroxypentynyl,
3-ethyl-3-hydroxypentyl,
3-ethyl-3-hydroxypentenyl,
3-ethyl-3-hydroxypentynyl,
3-ethyl-3-hydroxy-4-methylpentyl,
3-ethyl-3-hydroxy-4-methylpentenyl,
3-ethyl-3-hydroxy-4-methylpentynyl,
3-propyl-3-hydroxypentyl,
3-propyl-3-hydroxypentenyl,
3-propyl-3-hydroxypentynyl,
1-hydroxy-2-methyl-1-(methylethyl)propyl,

1-hydroxycycloalkenyl; and
1-hydroxycycloalkyl.


The term, "3-methyl-3-hydroxypentyl" refers to the radical having the structural formula:




The term, "3-methyl-3-hydroxypentenyl" refers to the radical having the structural formula (both cis and trans isomers):




The term, "3-methyl-3-hydroxypentynyl" refers to the radical having the structural formula:




The term, "3-ethyl-3-hydroxypentyl" refers to the radical having the structural formula:




The term, "3-ethyl-3-hydroxypentenyl" refers to the radical having the structural formula (both cis and trans isomers):




The term, "3-ethyl-3-hydroxypentynyl" refers to the radical having the structural formula:




The term, "3-propyl-3-hydroxypentyl" refers to the radical having the structural formula:




The term, "3-propyl-3-hydroxypentenyl" refers to the radical having the structural formula (both cis and trans isomers):




The term, "3-propyl-3-hydroxypentynyl" refers to the radical having the structural formula:




The term, "3-ethyl-3-hydroxy-4-methylpentyl" refers to the radical having the structural formula:




The term, "3-ethyl-3-hydroxy-4-methylpentenyl" refers to the radical having the structural formula (both cis and trans isomers):




The term, "3-ethyl-3-hydroxy-4-methylpentynyl" refers to the radical having the structural formula:




The term, "1-hydroxy-2-methyl-1-(methylethyl)propyl" refers to the radical having the structural formula:




The term, "3-methyl-3-hydroxy-4,4-dimethylpentyl" refers to the radical having the structural formula:




The term, "3-methyl-3-hydroxy-4,4-dimethylpentenyl." refers to the radical having the structural formula (both cis and trans isomers):




The term, "3-methyl-3-hydroxy-4,4-dimethylpentynyl" refers to the radical having the structural formula:




The term, "3-ethyl-3-hydroxy-4,4-dimethylpentynyl" refers to the radical having the structural formula:



The term, "3-ethyl-3-hydroxy-4,4-dimethylpentenyl" refers to the radical having the structural formula (both cis and trans isomers):




The term, "3-ethyl-3-hydroxy-4,4-dimethylpentynyl" refers to the radical having the structural formula:




The term, "1-hydroxycycloalkenyl" refers to a radical selected from 1-hydroxycyclopentenyl, 1-hydroxycyclohexenyl, 1-hydroxycycloheptenyl, or 1-hydroxycyclooctenyl.

The term "hydroxycycloalkyl" refers to a radical having the general structural formula:



where w is an integer from 1 to 6 and the hydroxyl radical is substituted on any ring carbon atom.

The term "1-hydroxycycloalkyl" refers to a radical having the general structural formula:



Examples of 1-hydroxycycloalkyl radicals are

1-hydroxycyclopropyl, 1-hydroxycyclobutyl, 1-hydroxycyclopentyl, 1-hydroxycyclohexyl, 1-hydroxycycloheptyl, and 1-hydroxycyclooctyl.

The abbreviation, "Me" means methyl.

The abbreviation, "Et" means ethyl.

The abbreviation, "iPr" means 1-methylethyl.

The abbreviation, "nPr" means n-propyl.

The abbreviation, "3Me3OH-Pentyl" means 3-methyl-3-hydroxypentyl.

The abbreviation, "3Me3OH-Pentenyl" means 3-methyl-3-hydroxypentenyl

The abbreviation, "3Me3OH-Pentynyl" means 3-methyl-3-hydroxypentynyl

The abbreviation, "3Et3OH-Pentyl" means 3-ethyl-3-hydroxypentyl.

The abbreviation, "3Et3OH-Pentenyl" means 3-ethyl-3-hydroxypentenyl

The abbreviation, "3Et3OH-Pentynyl" means 3-ethyl-3-hydroxypentynyl

The abbreviation, "3Pr3OH-Pentyl" means 3-propyl-3-hydroxypentyl.

The abbreviation, "3Pr3OH-Pentenyl" means 3-propyl-3-hydroxypentenyl.

The abbreviation, "3Pr3OH-Pentynyl" means 3-propyl-3-hydroxypentynyl.

The abbreviation, "3Et3OH4Me-Pentyl" means 3-ethyl-3-hydroxy-4-methylpentyl.


The abbreviation, "3Et3OH4Me-Pentenyl" means 3-ethyl-3-hydroxy-4-methylpentenyl,

The abbreviation, "3Et3OH4Me-Pentynyl" means 3-ethyl-3-hydroxy-4-methylpentynyl.

The abbreviation, "1OH2Me1MeEt-Propyl" means 1-hydroxy-2-methyl-1-(methylethyl)propyl.

Compounds of the Invention:

The compounds of the invention with vitamin receptor modulating (VDRM) activity are represented by formula (IA) or a pharmaceutically acceptable salt or a prodrug derivative thereof:



wherein





R and R' are independently C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, or together R and R' form a substituted or unsubstituted, saturated or unsaturated carbocyclic ring having from 3 to 8 carbon atoms;

RP₃ and RB are independently selected from the group consisting of hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, -O-C₁-C₅ alkyl, -S-C₁-C₅ alkyl, -O-C₁-C₅ fluoroalkyl, -CN, -NO₂, acetyl, -S-C₁-C₅ fluoroalkyl, C₂-C₅ alkenyl, C₃-C₅ cycloalkyl, and C₃-C₅ cycloalkenyl;

RP, RT₃, and RB' are independently selected from hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, -O-C₁-C₅ alkyl, -S-C₁-C₅ alkyl, -O-C₁-C₅ fluoroalkyl, -CN, -NO₂, acetyl, -S-C₁-C₅ fluoroalkyl, C₂-C₅ alkenyl, C₃-C₅ cycloalkyl, or C₃-C₅ cycloalkenyl;

(L_{P1}), (L_{P2}), and (L_{TB}) are divalent linking groups independently selected from the group consisting of

a bond





R40


R40



O



' and



where m is 0, 1, or 2, and each R40 is independently hydrogen, C₁-C₅ alkyl, or C₁-C₅ fluoroalkyl;

Z_p is

branched C₃-C₅ alkyl,
3-methyl-3-hydroxypentyl,
3-methyl-3-hydroxypentenyl,
3-methyl-3-hydroxypentynyl,
3-ethyl-3-hydroxypentyl,
3-ethyl-3-hydroxypentenyl,
3-ethyl-3-hydroxypentynyl,
3-ethyl-3-hydroxy-4-methylpentyl,
3-ethyl-3-hydroxy-4-methylpentenyl,
3-ethyl-3-hydroxy-4-methylpentynyl,
3-propyl-3-hydroxypentyl,
3-propyl-3-hydroxypentenyl,
3-propyl-3-hydroxypentynyl,
1-hydroxy-2-methyl-1-(methylethyl)propyl,
3-methyl-3-hydroxy-4,4-dimethylpentyl,
3-methyl-3-hydroxy-4,4-dimethylpentenyl,
3-methyl-3-hydroxy-4,4-dimethylpentyl,
3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
3-ethyl-3-hydroxy-4,4-dimethylpentenyl,
3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
1-hydroxycyclopentenyl,
1-hydroxycyclohexenyl,
1-hydroxycycloheptenyl,
1-hydroxycyclooctenyl,
1-hydroxycyclopropyl,
1-hydroxycyclobutyl,
1-hydroxycyclopentyl,
1-hydroxycyclohexyl,
1-hydroxycycloheptyl, or
1-hydroxycyclooctyl;

provided, however, that when

Z_p is

3-methyl-3-hydroxypentyl,
 3-methyl-3-hydroxypentenyl,
 3-methyl-3-hydroxypentynyl,
 3-ethyl-3-hydroxypentyl,
 3-ethyl-3-hydroxypentenyl,
 3-ethyl-3-hydroxypentynyl,
 3-ethyl-3-hydroxy-4-methylpentyl,
 3-ethyl-3-hydroxy-4-methylpentenyl,
 3-ethyl-3-hydroxy-4-methylpentynyl,
 3-propyl-3-hydroxypentyl,
 3-propyl-3-hydroxypentenyl,
 3-propyl-3-hydroxypentynyl,
 3-methyl-3-hydroxy-4,4-dimethylpentyl,
 3-methyl-3-hydroxy-4,4-dimethylpentenyl,
 3-methyl-3-hydroxy-4,4-dimethylpentynyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentenyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentynyl, or
 1-hydroxy-2-methyl-1-(methylethyl)propyl;

then (L_{P1}) and (L_{P2}) combine as a bond;

Z_{TB} is selected from

-O-(C₁-C₅ alkyl),
 -O-(C₂-C₅ alkenyl),
 -O-(C₃-C₅ cycloalkyl),
 -O-(C₃-C₅ cycloalkenyl),
 -O-(C₁-C₅ hydroxyalkyl),
 -O-(C₁-C₅ fluoroalkyl),
 -O-(C₁-C₅ alkyl)-phenyl,
 -O-(C₁-C₅ alkyl)-(O)-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl) NH₂,

-O-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl)₂ ,
-O-(C₁-C₅ alkyl)-C(O)-NH₂,
-O-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-C(O)-OH,
-O-(C₁-C₅ alkyl)-C(O)-NH-5-tetrazolyl,
-O-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-NH₂,
-O-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-O-(C₁-C₅ alkyl)-N-pyrrolidine,
-O-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-O-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-SO₂-NH₂,
-O-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-NH₂,
-O-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂ ,
-O-(C₁-C₅ alkyl)-5-tetrazolyl,
-O-CH₂-CO₂H,
-O-CH₂-5-tetrazolyl,

-O-(C₁-C₅ alkyl),
-O-C(O)-NH₂,
-O-C(O)-N-(CH₃)₂,
-O-C(S)-N-(CH₃)₂,
-O-C(O)-O-(C₁-C₅ alkyl),
-O-(5-tetrazolyl),
-O-SO₂-(C₁-C₅ alkyl),
-O-SO₂-NH₂,
-O-SO₂-NH-(C₁-C₅ alkyl),
-O-SO₂-N-(C₁-C₅ alkyl)₂,
-O-S(O)-(C₁-C₅ alkyl),
-O-S(O)-NH₂,
-O-S(O)-NH-(C₁-C₅ alkyl),
-O-S(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl),
-S-(C₂-C₅ alkenyl),
-S-(C₃-C₅ cycloalkyl),
-S-(C₃-C₅ cycloalkenyl),
-S-(C₁-C₅ fluoroalkyl),
-S-(C₁-C₅ hydroxyalkyl),
-S-(C₁-C₅ alkyl)-phenyl,
-S-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-OH,
-S-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-NH₂,
-S-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl) NH₂,

-S-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-S-(C₁-C₅ alkyl)-N-pyrrolidine,
-S-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-S-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-SO₂-NH₂,
-S-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-5-tetrazolyl,
-S-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-S(O)-NH₂,
-S-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₂-C₅ alkenyl),
-SO₂-(C₃-C₅ cycloalkyl),
-SO₂-(C₃-C₅ cycloalkenyl),
-SO₂-(C₁-C₅ hydroxyalkyl),
-SO₂-(C₁-C₅ fluoroalkyl),
-SO₂-(C₁-C₅)-phenyl,
-SO₂-NH₂,
-SO₂-NH-(C₁-C₅ alkyl),
-SO₂-NH-CH₂-C(O)OH,
-SO₂-NH-CH₂-C(O)(O-C₁-C₅ alkyl),

-SO₂-NH-(C₁-C₅ alkyl)-C(O)OH,
-SO₂-NH-(C₁-C₅ alkyl)-C(O)(O-C₁-C₅ alkyl),
-SO₂-NHC(O)-(C₃-C₆ cycloalkyl),
-SO₂-NH-C(O)-(C₁-C₅ alkyl),
-SO₂-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl) NH₂,
-SO₂-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-C(O)-NH₂,
-SO₂-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-SO₂-(C₁-C₅ alkyl)-N-pyrrolidine,
-SO₂-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-SO₂-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-OH,
-SO₂-(C₁-C₅ alkyl)-5-tetrazolyl,
-SO₂-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-SO₂-NH₂,
-SO₂-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₂-C₅ alkenyl),
-SO₂-(C₃-C₅ cycloalkyl),

-SO₂-(C₃-C₅ cycloalkenyl),
-SO₂-(C₁-C₅ hydroxyalkyl),
-SO₂-(C₁-C₅ fluoroalkyl),
-SO₂-(C₁-C₅)-phenyl,
-SO₂-N=CHN(C₁-C₅ alkyl)₂,
-S(O)-NH₂,
-S(O)-NH-(C₁-C₅ alkyl),
-S(O)-NH-CH₂-C(O)OH
-S(O)-NH-(C₁-C₅ alkyl)-C(O)OH,
-S(O)-NH-CH₂-C(O)(O-C₁-C₅ alkyl),
-S(O)-NH-(C₁-C₅ alkyl)-C(O)(O-C₁-C₅ alkyl),
-S(O)HC(O)-(C₃-C₆ cycloalkyl),
-S(O)-NH-C(O)-(C₁-C₅ alkyl),
-S(O)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-C(O)-NH₂,
-S(O)-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-NH-S(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-S(O)-(C₁-C₅ alkyl)-N-pyrrolidine,
-S(O)-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-S(O)-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-OH,

-S(O)-(C₁-C₅ alkyl)-5-tetrazolyl,
-S(O)-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-SO₂-NH₂,
-S(O)-(C₁-C₅ alkyl)-S(O)-NH₂,
-S(O)-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-S(O)-N=CHN(C₁-C₅ alkyl)₂,
-NHC(S)NH₂,
-NHC(S)NH-(C₁-C₅ alkyl),
-NHC(S)N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₂-C₅ alkenyl),
-NHC(S)NH-(C₃-C₅ cycloalkyl),
-NHC(S)NH-(C₃-C₅ cycloalkenyl),
-NHC(S)NH-(C₁-C₅ fluoroalkyl),
-NHC(S)NH-C₁-C₅ hydroxyalkyl,
-NHC(S)NH-(C₁-C₅ fluoroalkyl),
-NHC(S)NH-phenyl,
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-OH,
-NHC(S)NH-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),

-NHC(S)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-NH-S(O)-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-NHC(S)NH-(C₁-C₅ alkyl)-N-pyrrolidine,
-NHC(S)NH-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-
3-yl),
-NHC(S)NH-(C₁-C₅ alkyl)-5-tetrazolyl,
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-NHC(O)NH₂,
-NHC(O)NH-(C₁-C₅ alkyl),
-NHC(O)N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₂-C₅ alkenyl),
-NHC(O)NH-(C₃-C₅ cycloalkyl),
-NHC(O)NH-(C₃-C₅ cycloalkenyl),
-NHC(O)NH-(C₁-C₅ hydroxyalkyl),
-NHC(O)NH-(C₁-C₅ fluoroalkyl),
-NHC(O)NH-phenyl,

-NHC(O)NH-(C₁-C₅ alkyl)-NH₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-NH₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-C(O)-NH₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
 -NHC(O)NH-(C₁-C₅ alkyl)-N-pyrrolidine,
 -NHC(O)NH-(C₁-C₅ alkyl)-
 (1-methylpyrrolidin-2-one-3-yl),
 -NHC(O)NH-(C₁-C₅ alkyl)-C(O)-OH,
 -NHC(O)NH-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-5-tetrazolyl,
 -NHC(O)NH-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-SO₂-NH₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-P(O)-O-(C₁-C₅ alkyl)₂,
 -NH₂,
 -NH-(C₁-C₅ alkyl),
 -NH-CH₂-C(O)OH,
 -N-(C₁-C₅ alkyl)₂,
 -NH-C(O)-NH₂,

-NH-C(O)-NH-(C₁-C₅ alkyl),
-NH-C(O)-N-(C₁-C₅ alkyl)₂,
-NH-C(O)-(C₁-C₅ alkyl),
-NH-SO₂-(C₁-C₅ alkyl),
-NH-S(O)-(C₁-C₅ alkyl),
-N(CH₃)(OCH₃),
-N(OH)(CH₃),
-N-pyrrolidin-2-one,
-N-pyrrolidine,
-(1-methylpyrrolidin-2-one-3-yl),
-CO₂H,
-CO₂Me,
-CO₂Et,
-C(O)CH₂S(O)Me,
-C(O)CH₂S(O)Et,
-C(O)CH₂S(O)₂Me,
-C(O)CH₂S(O)₂Et,
-C(O)CH₂CH₂S(O)Me,
-C(O)CH₂CH₂S(O)Et,
-C(O)CH₂CH₂S(O)₂Me,
-C(O)CH₂CH₂S(O)₂Et,
-C(O)CH(Me)CH₂CO₂H,
-C(O)CH(Me)CH₂CO₂Me,
-C(O)CH(Me)CH₂CO₂Et,
-C(O)CH(Me)CH₂CO₂iPr,
-C(O)CH(Me)CH₂CO₂tBu,
-C(O)CH(Me)CH(Me)CO₂H,
-C(O)CH(Me)CH(Me)CO₂Me,
-C(O)CH(Me)CH(Me)CO₂Et,
-C(O)CH(Me)CH(Me)CO₂iPr,

-C(O)CH(Me)CH(Me)CO₂tBu,
-C(O)CH(Me)C(Me)₂CO₂H,
-C(O)CH(Me)C(Me)₂CO₂Me,
-C(O)CH(Me)C(Me)₂CO₂Et,
-C(O)CH(Me)C(Me)₂CO₂iPr,
-C(O)CH(Me)C(Me)₂CO₂tBu,
-C(O)CH(Me)CH(Et)CO₂H,
-C(O)CH(Me)CH(Et)CO₂Me,
-C(O)CH(Me)CH(Et)CO₂Et,
-C(O)CH(Me)CH(Et)CO₂iPr,
-C(O)CH(Me)CH(Et)CO₂tBu,
-C(O)C(O)OH,
-C(O)C(O)NH₂,
-C(O)C(O)NHMe,
-C(O)C(O)NMe₂,
-C(O)NH₂,
-C(O)NMe₂,
-C(O)NH-CH₂-C(O)OH,
-C(O)NH-CH₂-C(O)OMe,
-C(O)NH-CH₂-C(O)OEt,
-C(O)NH-CH₂-C(O)OiPr,
-C(O)NH-CH₂-C(O)OtBu,
-C(O)NH-CH(Me)-C(O)OH,
-C(O)NH-CH(Me)-C(O)OMe,
-C(O)NH-CH(Me)-C(O)OEt,
-C(O)NH-CH(Me)-C(O)iPr,
-C(O)NH-CH(Me)-C(O)tBu,
-C(O)NH-CH(Et)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OMe,
-C(O)NH-C(Me)₂-C(O)OEt,

-C(O)NH-C(Me)₂-C(O)iPr,
-C(O)NH-C(Me)₂-C(O)tBu,
-C(O)NH-CMe(Et)-C(O)OH,
-C(O)NH-CH(F)-C(O)OH,
-C(O)NH-CH(CF₃)-C(O)OH,
-C(O)NH-CH(OH)-C(O)OH,
-C(O)NH-CH(cyclopropyl)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-CF(Me)-C(O)OH,
-C(O)NH-C(Me)(CF₃)-C(O)OH,
-C(O)NH-C(Me)(OH)-C(O)OH,
-C(O)NH-C(Me)(cyclopropyl)CO₂H
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH₂-C(OMe),
-C(O)NMe-CH₂-C(O)OEt,
-C(O)NMe-CH₂-C(O)OiPr,
-C(O)NMe-CH₂-C(O)tBu,
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH(Me)-C(O)OH,
-C(O)NMe-CH(F)-C(O)OH,
-C(O)NMe-CH(CF₃)-C(O)OH,
-C(O)NMe-CH(OH)-C(O)OH,
-C(O)NMe-CH(cyclopropyl)-C(O)OH,
-C(O)NMe-C(Me)₂-C(O)OH,
-C(O)NMe-CF(Me)-C(O)OH,
-C(O)NMe-C(Me)(CF₃)-C(O)OH,
-C(O)NMe-C(Me)(OH)-C(O)OH,
-C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,
-C(O)NHS(O)Me,
-C(O)NHSO₂Me,
-C(O)-NH-5-tetrazolyl,
-C(O)NHS(O)Me,

-C(O)NHS(O)Et,
-C(O)NH₂SO₂Me,
-C(O)NH₂SO₂Et,
-C(O)NHS(O)iPr,
-C(O)NH₂SO₂iPr,
-C(O)NHS(O)tBu,
-C(O)NH₂SO₂tBu,
-C(O)NHCH₂S(O)Me,
-C(O)NHCH₂S(O)Et,
-C(O)NHCH₂SO₂Me,
-C(O)NHCH₂SO₂Et,
-C(O)NHCH₂CH₂S(O)Me,
-C(O)NHCH₂CH₂S(O)Et,
-C(O)NHCH₂CH₂SO₂Me,
-C(O)NHCH₂CH₂SO₂Et,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)SO₂Me,
-C(O)-N(Me)-5-tetrazolyl,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)S(O)Et,
-C(O)N(Me)SO₂Me,
-C(O)N(Me)SO₂Et,
-C(O)N(Me)S(O)iPr,
-C(O)N(Me)SO₂iPr,
-C(O)N(Me)S(O)tBu,
-C(O)N(Me)SO₂tBu,
-C(O)N(Me)CH₂S(O)Me,
-C(O)N(Me)CH₂S(O)Et,
-C(O)N(Me)CH₂SO₂Me,
-C(O)N(Me)CH₂SO₂Et,

-C(O)N(Me)CH₂CH₂S(O)Me,
-C(O)N(Me)CH₂CH₂S(O)Et,
-C(O)N(Me)CH₂CH₂SO₂Me,
-C(O)N(Me)CH₂CH₂SO₂Et,
-CH₂CO₂H,
-CH₂-5-tetrazolyl,
-CH₂CO₂Me,
-CH₂CO₂Et,
-CH₂NHS(O)Me,
-CH₂NHS(O)Et,
-CH₂NHSO₂Me,
-CH₂NHSO₂Et,
-CH₂NHS(O)iPr,
-CH₂NHSO₂iPr,
-CH₂NHS(O)tBu,
-CH₂NHSO₂tBu,
-CH₂NHCH₂CH₂SO₂CH₃,
-CH₂NH(CH₂CO₂H),
-CH₂N(C(O)Me)(CH₂CO₂H),
-CH₂-N-pyrrolidin-2-one,
-CH₂-(1-methylpyrrolidin-2-one-3-yl),
-CH₂S(O)Me,
-CH₂S(O)Et,
-CH₂S(O)₂Me,
-CH₂S(O)₂Et,
-CH₂S(O)iPr,
-CH₂S(O)₂iPr,
-CH₂S(O)tBu,
-CH₂S(O)₂tBu,

-CH₂CO₂H, CH₂C(O)NH₂,
-CH₂C(O)NMe₂,
-CH₂C(O)NHMe,
-CH₂C(O)-N-pyrrolidine,
-CH₂S(O)₂Me, CH₂S(O)Me,
-CH(OH)CO₂H,
-CH(OH)C(O)NH₂,
-CH(OH)C(O)NHMe,
-CH(OH)C(O)NMe₂,
-CH(OH)C(O)NEt₂,
-CH₂CH₂CO₂H,
-CH₂CH₂CO₂Me,
-CH₂CH₂CO₂Et,
-CH₂CH₂C(O)NH₂,
-CH₂CH₂C(O)NHMe,
-CH₂CH₂C(O)NMe₂,
-CH₂CH₂-5-tetrazolyl,
-CH₂CH₂S(O)₂Me,
-CH₂CH₂S(O)Me,
-CH₂CH₂S(O)₂Et,
-CH₂CH₂S(O)Et,
-CH₂CH₂S(O)iPr,
-CH₂CH₂S(O)₂iPr,
-CH₂CH₂S(O)tBu,
-CH₂CH₂S(O)₂tBu,
-CH₂CH₂S(O)NH₂,
-CH₂CH₂S(O)NHMe,
-CH₂CH₂S(O)NMe₂,
-CH₂CH₂S(O)₂NH₂,

-CH₂CH₂S(O)₂NHMe

-CH₂CH₂S(O)₂NMe₂,

-CH₂CH₂CH₂S(O)Me,






-CH₂CH₂CH₂S(O)Et,









-CH₂CH₂CH₂S(O)₂Me,


-CH₂CH₂CH₂S(O)₂Et,

-C(O)OH,

-5-tetrazolyl,







-1,3,4-oxadiazolin-2-one-5-yl,
 -imidazolidine-2,4-dione-5-yl,
 -isoxazol-3-ol-yl, or
 -1,3,4-oxadiazolin-2-thione-5-yl;

provided that RB is substituted at either the 6 or 7 position of the benzothiophene ring, except that RB is substituted only at the 7 position of the benzothiophene ring when Z_{TB} is at the 6 position.; and


provided that -(L_{TB})-Z_{TB} is substituted at either the 5 or 6 position of the benzothiophene ring; and

provided that RB is substituted at either the 6 or 7 position of the benzothiophene ring, except that RB is substituted only at the 7 position of the benzothiophene ring when the group $-(L_{TB})-Z_{TB}$ is at the 6 position.; and

provided that RB' is substituted at either the 4 or 5 position of the benzothiophene ring, except that RB' is substituted only at the 5 position of the benzothiophene ring when the group $-(L_{TB})-Z_{TB}$ is at the 6 position of the phenyl ring; and

provided that RP is substituted at either the 2, or 5 or 6 position of the phenyl ring.

The compounds of the invention with vitamin receptor modulating (VDRM) activity are represented by formula (IB) or a pharmaceutically acceptable salt or a prodrug derivative thereof:



wherein the variables R, R', RP, RP₃, Z_p, RB₇, RB, RB₄, RT₃, and Z_{BT} are as hereinafter defined.

wherein

R and R' are independently C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, or together R and R' form a substituted or unsubstituted, saturated or unsaturated carbocyclic ring having from 3 to 8 carbon atoms;





RP, RB₄, RT₃ and RB are independently selected from the group consisting of hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, -O-C₁-C₅ alkyl, -S-C₁-C₅ alkyl, -O-C₁-C₅ fluoroalkyl, -CN, -NO₂, acetyl, -S-C₁-C₅ fluoroalkyl, C₂-C₅ alkenyl, C₃-C₅ cycloalkyl, and C₃-C₅ cycloalkenyl;

RP₃ and RB₇ are independently selected from hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, -O-C₁-C₅ alkyl, -S-C₁-C₅ alkyl, -O-C₁-C₅ fluoroalkyl, -CN, -

NO₂, acetyl, -S-C₁-C₅ fluoroalkyl, C₂-C₅ alkenyl, C₃-C₅ cycloalkyl, or C₃-C₅ cycloalkenyl;


(L_{P1}), (L_{P2}), and (L_{BT}) are divalent linking groups independently selected from the group consisting of

a bond





R40


R40



R40



and



where m is 0, 1, or 2, and each R40 is independently hydrogen, C₁-C₅ alkyl, or C₁-C₅ fluoroalkyl;

Z_P is

branched C₃-C₅ alkyl,

3-methyl-3-hydroxypentyl,

3-methyl-3-hydroxypentenyl,

3-methyl-3-hydroxypentynyl,

3-ethyl-3-hydroxypentyl,
3-ethyl-3-hydroxypentenyl,
3-ethyl-3-hydroxypentynyl,
3-ethyl-3-hydroxy-4-methylpentyl,
3-ethyl-3-hydroxy-4-methylpentenyl,
3-ethyl-3-hydroxy-4-methylpentynyl,
3-propyl-3-hydroxypentyl,
3-propyl-3-hydroxypentenyl,
3-propyl-3-hydroxypentynyl,
1-hydroxy-2-methyl-1-(methylethyl)propyl,
3-methyl-3-hydroxy-4,4-dimethylpentyl,
3-methyl-3-hydroxy-4,4-dimethylpentenyl,
3-methyl-3-hydroxy-4,4-dimethylpentyl,
3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
3-ethyl-3-hydroxy-4,4-dimethylpentenyl,
3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
1-hydroxycyclopentenyl,
1-hydroxycyclohexenyl,
1-hydroxycycloheptenyl,
1-hydroxycyclooctenyl,
1-hydroxycyclopropyl,
1-hydroxycyclobutyl,
1-hydroxycyclopentyl,
1-hydroxycyclohexyl,
1-hydroxycycloheptyl, or
1-hydroxycyclooctyl;

provided, however, that when

Z_p is

3-methyl-3-hydroxypentyl,
3-methyl-3-hydroxypentenyl,
3-methyl-3-hydroxypentynyl,
3-ethyl-3-hydroxypentyl,
3-ethyl-3-hydroxypentenyl,
3-ethyl-3-hydroxypentynyl,

3-ethyl-3-hydroxy-4-methylpentyl,
 3-ethyl-3-hydroxy-4-methylpentenyl,
 3-ethyl-3-hydroxy-4-methylpentynyl,
 3-propyl-3-hydroxypentyl,
 3-propyl-3-hydroxypentenyl,
 3-propyl-3-hydroxypentynyl,
 3-methyl-3-hydroxy-4,4-dimethylpentyl,
 3-methyl-3-hydroxy-4,4-dimethylpentenyl,
 3-methyl-3-hydroxy-4,4-dimethylpentyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentenyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentynyl, or
 1-hydroxy-2-methyl-1-(methylethyl)propyl;

then (L_{P1}) and (L_{P2}) combine as a bond;

Z_{BT} is selected from

-O-(C₁-C₅ alkyl),
 -O-(C₂-C₅ alkenyl),
 -O-(C₃-C₅ cycloalkyl),
 -O-(C₃-C₅ cycloalkenyl),
 -O-(C₁-C₅ hydroxyalkyl),
 -O-(C₁-C₅ fluoroalkyl),
 -O-(C₁-C₅ alkyl)-phenyl,
 -O-(C₁-C₅ alkyl)-(O)-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl) NH₂,
 -O-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl)₂,
 -O-(C₁-C₅ alkyl)-C(O)-NH₂,
 -O-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
 -O-(C₁-C₅ alkyl)-C(O)-OH,
 -O-(C₁-C₅ alkyl)-C(O)-NH-5-tetrazolyl,
 -O-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),

-O-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-NH₂,
-O-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-O-(C₁-C₅ alkyl)-N-pyrrolidine,
-O-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-O-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-SO₂-NH₂,
-O-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-NH₂,
-O-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-5-tetrazolyl,
-O-CH₂-CO₂H,
-O-CH₂-5-tetrazolyl,
-O-(C₁-C₅ alkyl),
-O-C(O)-NH₂,
-O-C(O)-N-(CH₃)₂,
-O-C(S)-N-(CH₃)₂,
-O-C(O)-O-(C₁-C₅ alkyl),
-O-(5-tetrazolyl),
-O-SO₂-(C₁-C₅ alkyl),
-O-SO₂-NH₂,

-O-SO₂-NH-(C₁-C₅ alkyl),
-O-SO₂-N-(C₁-C₅ alkyl)₂,
-O-S(O)-(C₁-C₅ alkyl),
-O-S(O)-NH₂,
-O-S(O)-NH-(C₁-C₅ alkyl),
-O-S(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl),
-S-(C₂-C₅ alkenyl),
-S-(C₃-C₅ cycloalkyl),
-S-(C₃-C₅ cycloalkenyl),
-S-(C₁-C₅ fluoroalkyl),
-S-(C₁-C₅ hydroxyalkyl),
-S-(C₁-C₅ alkyl)-phenyl,
-S-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-OH,
-S-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-NH₂,
-S-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl) NH₂,
-S-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-S-(C₁-C₅ alkyl)-N-pyrrolidine,
-S-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-S-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-SO₂-NH₂,

-S-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-5-tetrazolyl,
-S-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-S(O)-NH₂,
-S-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₂-C₅ alkenyl),
-SO₂-(C₃-C₅ cycloalkyl),
-SO₂-(C₃-C₅ cycloalkenyl),
-SO₂-(C₁-C₅ hydroxyalkyl),
-SO₂-(C₁-C₅ fluoroalkyl),
-SO₂-(C₁-C₅)-phenyl,
-SO₂-NH₂,
-SO₂-NH-(C₁-C₅ alkyl),
-SO₂-NH-CH₂-C(O)OH,
-SO₂-NH-CH₂-C(O)(O-C₁-C₅ alkyl),
-SO₂-NH-(C₁-C₅ alkyl)-C(O)OH,
-SO₂-NH-(C₁-C₅ alkyl)-C(O)(O-C₁-C₅ alkyl),
-SO₂-NHC(O)-(C₃-C₆ cycloalkyl),
-SO₂-NH-C(O)-(C₁-C₅ alkyl),
-SO₂-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl) NH₂,

-SO₂-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-C(O)-NH₂,
-SO₂-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-SO₂-(C₁-C₅ alkyl)-N-pyrrolidine,
-SO₂-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-SO₂-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-OH,
-SO₂-(C₁-C₅ alkyl)-5-tetrazolyl,
-SO₂-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-SO₂-NH₂,
-SO₂-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₂-C₅ alkenyl),
-SO₂-(C₃-C₅ cycloalkyl),
-SO₂-(C₃-C₅ cycloalkenyl),
-SO₂-(C₁-C₅ hydroxyalkyl),
-SO₂-(C₁-C₅ fluoroalkyl),
-SO₂-(C₁-C₅)-phenyl,
-SO₂-N=CHN(C₁-C₅ alkyl)₂,
-S(O)-NH₂,
-S(O)-NH-(C₁-C₅ alkyl),
-S(O)-NH-CH₂-C(O)OH

-S(O)-NH-(C₁-C₅ alkyl)-C(O)OH,
-S(O)-NH-CH₂-C(O)(O-C₁-C₅ alkyl),
-S(O)-NH-(C₁-C₅ alkyl)-C(O)(O-C₁-C₅ alkyl),
-S(O)HC(O)-(C₃-C₆ cycloalkyl),
-S(O)-NH-C(O)-(C₁-C₅ alkyl),
-S(O)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-C(O)-NH₂,
-S(O)-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-NH-S(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-S(O)-(C₁-C₅ alkyl)-N-pyrrolidine,
-S(O)-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-S(O)-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-OH,
-S(O)-(C₁-C₅ alkyl)-5-tetrazolyl,
-S(O)-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-SO₂-NH₂,
-S(O)-(C₁-C₅ alkyl)-S(O)-NH₂,
-S(O)-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,

-S(O)-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
 -S(O)-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
 -S(O)-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
 -S(O)-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
 -S(O)-N=CHN(C₁-C₅ alkyl)₂,
 -NHC(S)NH₂,
 -NHC(S)NH-(C₁-C₅ alkyl),
 -NHC(S)N-(C₁-C₅ alkyl)₂,
 -NHC(S)NH-(C₂-C₅ alkenyl),
 -NHC(S)NH-(C₃-C₅ cycloalkyl),
 -NHC(S)NH-(C₃-C₅ cycloalkenyl),
 -NHC(S)NH-(C₁-C₅ fluoroalkyl),
 -NHC(S)NH-C₁-C₅ hydroxyalkyl,
 -NHC(S)NH-(C₁-C₅ fluoroalkyl),
 -NHC(S)NH-phenyl,
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-OH,
 -NHC(S)NH-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-NH₂,
 -NHC(S)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-NH₂,
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
 -NHC(S)NH-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-NH-S(O)-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
 -NHC(S)NH-(C₁-C₅ alkyl)-N-pyrrolidine,

-NHC(S)NH-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-NHC(S)NH-(C₁-C₅ alkyl)-5-tetrazolyl,
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-NHC(O)NH₂,
-NHC(O)NH-(C₁-C₅ alkyl),
-NHC(O)N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₂-C₅ alkenyl),
-NHC(O)NH-(C₃-C₅ cycloalkyl),
-NHC(O)NH-(C₃-C₅ cycloalkenyl),
-NHC(O)NH-(C₁-C₅ hydroxyalkyl),
-NHC(O)NH-(C₁-C₅ fluoroalkyl),
-NHC(O)NH-phenyl,
-NHC(O)NH-(C₁-C₅ alkyl)-NH₂,
-NHC(O)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-NH₂,
-NHC(O)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-NH₂,

-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-NHC(O)NH-(C₁-C₅ alkyl)-N-pyrrolidine,
-NHC(O)NH-(C₁-C₅ alkyl)-
 (1-methylpyrrolidin-2-one-3-yl),
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-OH,
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-5-tetrazolyl,
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-NH₂,
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₁-C₅ alkyl)-P(O)-O-(C₁-C₅ alkyl)₂ ,
-NH₂,
-NH-(C₁-C₅ alkyl),
-NH-CH₂-C(O)OH,
-N-(C₁-C₅ alkyl)₂,
-NH-C(O)-NH₂,
-NH-C(O)-NH-(C₁-C₅ alkyl),
-NH-C(O)-N-(C₁-C₅ alkyl)₂,
-NH-C(O)-(C₁-C₅ alkyl),
-NH-SO₂-(C₁-C₅ alkyl),
-NH-S(O)-(C₁-C₅ alkyl),
-N(CH₃)(OCH₃),
-N(OH)(CH₃),
-N-pyrrolidin-2-one,
-N-pyrrolidine,

-(1-methylpyrrolidin-2-one-3-yl),
-CO₂H,
-CO₂Me,
-CO₂Et,
-C(O)CH₂S(O)Me,
-C(O)CH₂S(O)Et,
-C(O)CH₂S(O)₂Me,
-C(O)CH₂S(O)₂Et,
-C(O)CH₂CH₂S(O)Me,
-C(O)CH₂CH₂S(O)Et,
-C(O)CH₂CH₂S(O)₂Me,
-C(O)CH₂CH₂S(O)₂Et,
-C(O)CH(Me)CH₂CO₂H,
-C(O)CH(Me)CH₂CO₂Me,
-C(O)CH(Me)CH₂CO₂Et,
-C(O)CH(Me)CH₂CO₂iPr,
-C(O)CH(Me)CH₂CO₂tBu,
-C(O)CH(Me)CH(Me)CO₂H,
-C(O)CH(Me)CH(Me)CO₂Me,
-C(O)CH(Me)CH(Me)CO₂Et,
-C(O)CH(Me)CH(Me)CO₂iPr,
-C(O)CH(Me)CH(Me)CO₂tBu,
-C(O)CH(Me)C(Me)₂CO₂H,
-C(O)CH(Me)C(Me)₂CO₂Me,
-C(O)CH(Me)C(Me)₂CO₂Et,
-C(O)CH(Me)C(Me)₂CO₂iPr,
-C(O)CH(Me)C(Me)₂CO₂tBu,
-C(O)CH(Me)CH(Et)CO₂H,
-C(O)CH(Me)CH(Et)CO₂Me,










-C(O)CH(Me)CH(Et)CO₂Et,
-C(O)CH(Me)CH(Et)CO₂iPr,
-C(O)CH(Me)CH(Et)CO₂tBu,
-C(O)C(O)OH,
-C(O)C(O)NH₂,
-C(O)C(O)NHMe,
-C(O)C(O)NMe₂,
-C(O)NH₂,
-C(O)NMe₂,
-C(O)NH-CH₂-C(O)OH,
-C(O)NH-CH₂-C(O)OMe,
-C(O)NH-CH₂-C(O)OEt,
-C(O)NH-CH₂-C(O)OiPr,
-C(O)NH-CH₂-C(O)OtBu,
-C(O)NH-CH(Me)-C(O)OH,
-C(O)NH-CH(Me)-C(O)OMe,
-C(O)NH-CH(Me)-C(O)OEt,
-C(O)NH-CH(Me)-C(O)iPr,
-C(O)NH-CH(Me)-C(O)tBu,
-C(O)NH-CH(Et)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OMe,
-C(O)NH-C(Me)₂-C(O)OEt,
-C(O)NH-C(Me)₂-C(O)iPr,
-C(O)NH-C(Me)₂-C(O)tBu,
-C(O)NH-CMe(Et)-C(O)OH,
-C(O)NH-CH(F)-C(O)OH,
-C(O)NH-CH(CF₃)-C(O)OH,
-C(O)NH-CH(OH)-C(O)OH,
-C(O)NH-CH(cyclopropyl)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-CF(Me)-C(O)OH,
















-C(O)NH-C(Me)(CF₃)-C(O)OH,
-C(O)NH-C(Me)(OH)-C(O)OH,
-C(O)NH-C(Me)(cyclopropyl)CO₂H
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH₂-C(O)OMe,
-C(O)NMe-CH₂-C(O)OEt,
-C(O)NMe-CH₂-C(O)OiPr,
-C(O)NMe-CH₂-C(O)tBu,
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH(Me)-C(O)OH,
-C(O)NMe-CH(F)-C(O)OH,
-C(O)NMe-CH(CF₃)-C(O)OH,
-C(O)NMe-CH(OH)-C(O)OH,
-C(O)NMe-CH(cyclopropyl)-C(O)OH,
-C(O)NMe-C(Me)₂-C(O)OH,
-C(O)NMe-CF(Me)-C(O)OH,
-C(O)NMe-C(Me)(CF₃)-C(O)OH,
-C(O)NMe-C(Me)(OH)-C(O)OH,
-C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,
-C(O)NHS(O)Me,
-C(O)NHSO₂Me,
-C(O)-NH-5-tetrazolyl,
-C(O)NHS(O)Me,
-C(O)NHS(O)Et,
-C(O)NHSO₂Me,
-C(O)NHSO₂Et,
-C(O)NHS(O)iPr,
-C(O)NHSO₂iPr,
-C(O)NHS(O)tBu,
-C(O)NHSO₂tBu,
-C(O)NHCH₂S(O)Me,
-C(O)NHCH₂S(O)Et,

-C(O)NHCH₂SO₂Me,
-C(O)NHCH₂SO₂Et,
-C(O)NHCH₂CH₂S(O)Me,
-C(O)NHCH₂CH₂S(O)Et,
-C(O)NHCH₂CH₂SO₂Me,
-C(O)NHCH₂CH₂SO₂Et,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)SO₂Me,
-C(O)-N(Me)-5-tetrazolyl,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)S(O)Et,
-C(O)N(Me)SO₂Me,
-C(O)N(Me)SO₂Et,
-C(O)N(Me)S(O)iPr,
-C(O)N(Me))SO₂iPr,
-C(O)N(Me))S(O)tBu,
-C(O)N(Me)SO₂tBu,
-C(O)N(Me)CH₂S(O)Me,
-C(O)N(Me)CH₂S(O)Et,
-C(O)N(Me)CH₂SO₂Me,
-C(O)N(Me)CH₂SO₂Et,
-C(O)N(Me)CH₂CH₂S(O)Me,
-C(O)N(Me)CH₂CH₂S(O)Et,
-C(O)N(Me)CH₂CH₂SO₂Me,
-C(O)N(Me)CH₂CH₂SO₂Et,
-CH₂CO₂H,
-CH₂-5-tetrazolyl,
-CH₂CO₂Me,

-CH₂CO₂Et,
-CH₂NHS(O)Me,
-CH₂NHS(O)Et,
-CH₂NHSO₂Me,
-CH₂NHSO₂Et,
-CH₂NHS(O)iPr,
-CH₂NHSO₂iPr,
-CH₂NHS(O)tBu,
-CH₂NHSO₂tBu,
-CH₂NHCH₂CH₂SO₂CH₃,
-CH₂NH(CH₂CO₂H),
-CH₂N(C(O)Me)(CH₂CO₂H),
-CH₂-N-pyrrolidin-2-one,
-CH₂-(1-methylpyrrolidin-2-one-3-yl),
-CH₂S(O)Me,
-CH₂S(O)Et,
-CH₂S(O)₂Me,
-CH₂S(O)₂Et,
-CH₂S(O)iPr,
-CH₂S(O)₂iPr,
-CH₂S(O)tBu,
-CH₂S(O)₂tBu,
-CH₂CO₂H, CH₂C(O)NH₂,
-CH₂C(O)NMe₂,
-CH₂C(O)NHMe,
-CH₂C(O)-N-pyrrolidine,
-CH₂S(O)₂Me, CH₂S(O)Me,
-CH(OH)CO₂H,
-CH(OH)C(O)NH₂,

-CH(OH)C(O)NHMe,
-CH(OH)C(O)NMe₂,
-CH(OH)C(O)NEt₂,
-CH₂CH₂CO₂H,
-CH₂CH₂CO₂Me,
-CH₂CH₂CO₂Et,
-CH₂CH₂C(O)NH₂,
-CH₂CH₂C(O)NHMe,
-CH₂CH₂C(O)NMe₂,
-CH₂CH₂-5-tetrazolyl,
-CH₂CH₂S(O)₂Me,
-CH₂CH₂S(O)Me,
-CH₂CH₂S(O)₂Et,
-CH₂CH₂S(O) Et,
-CH₂CH₂S(O)iPr,
-CH₂CH₂S(O)₂iPr,
-CH₂CH₂S(O)tBu,
-CH₂CH₂S(O)₂tBu,
-CH₂CH₂S(O)NH₂,
-CH₂CH₂S(O)NHMe,
-CH₂CH₂S(O)NMe₂,
-CH₂CH₂S(O)₂NH₂,
-CH₂CH₂S(O)₂NHMe,
-CH₂CH₂S(O)₂NMe₂,
-CH₂CH₂CH₂S(O)Me,
-CH₂CH₂CH₂S(O)Et,
-CH₂CH₂CH₂S(O)₂Me,
-CH₂CH₂CH₂S(O)₂Et,
-C(O)OH,
-5-tetrazolyl,





-1,3,4-oxadiazolin-2-one-5-yl,

-imidazolidine-2,4-dione-5-yl,
 -isoxazol-3-ol-yl, or
 -1,3,4-oxadiazolin-2-thione-5-yl;

provided that RP is substituted at either the 2, 5, or 6 position of the phenyl ring.

Preferred embodiments of the Invention:

The compound of Formula IA having as preferred substituents;

R and R' are independently methyl or ethyl;

RP and RT₃ are independently, hydrogen or methyl;

RP₃ and RB are independently hydrogen, methyl, ethyl, -O-methyl, or cyclopropyl;

(L_{P1}) and (L_{TB}) divalent linking groups are both bonds;

(L_{P2}) is a bond, -CH₂-, -CH(OH)-, or -C(Me)OH-;

Z_P is 1,1-dimethylethyl; 1-hydroxycyclopentyl, 1-hydroxycyclohexyl, 3-ethyl-3-hydroxypentyl, 3-ethyl-3-hydroxypentenyl, 3-ethyl-3-hydroxypentynyl;

Z_{TB} is

-CO₂H,
 -CO₂Me,
 -CO₂Et,
 -C(O)CH₂S(O)Me,
 -C(O)CH₂S(O)Et,
 -C(O)CH₂S(O)₂Me,
 -C(O)CH₂S(O)₂Et,
 -C(O)CH₂CH₂S(O)Me,
 -C(O)CH₂CH₂S(O)Et,
 -C(O)CH₂CH₂S(O)₂Me,
 -C(O)CH₂CH₂S(O)₂Et,
 -C(O)CH(Me)CH₂CO₂H,
 -C(O)CH(Me)CH₂CO₂Me,

-C(O)CH(Me)CH₂CO₂Et,
-C(O)CH(Me)CH₂CO₂iPr,
-C(O)CH(Me)CH₂CO₂tBu,
-C(O)CH(Me)CH(Me)CO₂H,
-C(O)CH(Me)CH(Me)CO₂Me,
-C(O)CH(Me)CH(Me)CO₂Et,
-C(O)CH(Me)CH(Me)CO₂iPr,
-C(O)CH(Me)CH(Me)CO₂tBu,
-C(O)CH(Me)C(Me)₂CO₂H,
-C(O)CH(Me)C(Me)₂CO₂Me,
-C(O)CH(Me)C(Me)₂CO₂Et,
-C(O)CH(Me)C(Me)₂CO₂iPr,
-C(O)CH(Me)C(Me)₂CO₂tBu,
-C(O)CH(Me)CH(Et)CO₂H,
-C(O)CH(Me)CH(Et)CO₂Me,
-C(O)CH(Me)CH(Et)CO₂Et,
-C(O)CH(Me)CH(Et)CO₂iPr,
-C(O)CH(Me)CH(Et)CO₂tBu,
-C(O)C(O)OH,
-C(O)C(O)NH₂,
-C(O)C(O)NHMe,
-C(O)C(O)NMe₂,
-C(O)NH₂,
-C(O)NMe₂,
-C(O)NH-CH₂-C(O)OH,
-C(O)NH-CH₂-C(O)OMe,
-C(O)NH-CH₂-C(O)OEt,
-C(O)NH-CH₂-C(O)OiPr,
-C(O)NH-CH₂-C(O)OtBu,
-C(O)NH-CH(Me)-C(O)OH,

-C(O)NH-CH(Me)-C(O)OMe,
-C(O)NH-CH(Me)-C(O)OEt,
-C(O)NH-CH(Me)-C(O)iPr,
-C(O)NH-CH(Me)-C(O)tBu,
-C(O)NH-CH(Et)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OMe,
-C(O)NH-C(Me)₂-C(O)OEt,
-C(O)NH-C(Me)₂-C(O)iPr,
-C(O)NH-C(Me)₂-C(O)tBu,
-C(O)NH-CMe(Et)-C(O)OH,
-C(O)NH-CH(F)-C(O)OH,
-C(O)NH-CH(CF₃)-C(O)OH,
-C(O)NH-CH(OH)-C(O)OH,
-C(O)NH-CH(cyclopropyl)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-CF(Me)-C(O)OH,
-C(O)NH-C(Me)(CF₃)-C(O)OH,
-C(O)NH-C(Me)(OH)-C(O)OH,
-C(O)NH-C(Me)(cyclopropyl)CO₂H
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH₂-C(O)OMe,
-C(O)NMe-CH₂-C(O)OEt,
-C(O)NMe-CH₂-C(O)OiPr,
-C(O)NMe-CH₂-C(O)tBu,
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH(Me)-C(O)OH,
-C(O)NMe-CH(F)-C(O)OH,
-C(O)NMe-CH(CF₃)-C(O)OH,
-C(O)NMe-CH(OH)-C(O)OH,
-C(O)NMe-CH(cyclopropyl)-C(O)OH,
-C(O)NMe-C(Me)₂-C(O)OH,

-C(O)NMe-CF(Me)-C(O)OH,
-C(O)NMe-C(Me)(CF₃)-C(O)OH,
-C(O)NMe-C(Me)(OH)-C(O)OH,
-C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,
-C(O)NHS(O)Me,
-C(O)NHSO₂Me,
-C(O)-NH-5-tetrazolyl,
-C(O)NHS(O)Me,
-C(O)NHS(O)Et,
-C(O)NHSO₂Me,
-C(O)NHSO₂Et,
-C(O)NHS(O)iPr,
-C(O)NHSO₂iPr,
-C(O)NHS(O)tBu,
-C(O)NHSO₂tBu,
-C(O)NHCH₂S(O)Me,
-C(O)NHCH₂S(O)Et,
-C(O)NHCH₂SO₂Me,
-C(O)NHCH₂SO₂Et,
-C(O)NHCH₂CH₂S(O)Me,
-C(O)NHCH₂CH₂S(O)Et,
-C(O)NHCH₂CH₂SO₂Me,
-C(O)NHCH₂CH₂SO₂Et,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)SO₂Me,
-C(O)-N(Me)-5-tetrazolyl,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)S(O)Et,
-C(O)N(Me)SO₂Me,
-C(O)N(Me)SO₂Et,
-C(O)N(Me)S(O)iPr,

-C(O)N(Me))SO₂iPr,
-C(O)N(Me))S(O)tBu,
-C(O)N(Me)SO₂tBu,
-C(O)N(Me)CH₂S(O)Me,
-C(O)N(Me)CH₂S(O)Et,
-C(O)N(Me)CH₂SO₂Me,
-C(O)N(Me)CH₂SO₂Et,
-C(O)N(Me)CH₂CH₂S(O)Me,
-C(O)N(Me)CH₂CH₂S(O)Et,
-C(O)N(Me)CH₂CH₂SO₂Me,
-C(O)N(Me)CH₂CH₂SO₂Et,
-CH₂CO₂H,
-CH₂-5-tetrazolyl,
-CH₂CO₂Me,
-CH₂CO₂Et,
-CH₂NHS(O)Me,
-CH₂NHS(O)Et,
-CH₂NHSO₂Me,
-CH₂NHSO₂Et,
-CH₂NHS(O)iPr,
-CH₂NHSO₂iPr,
-CH₂NHS(O)tBu,
-CH₂NHSO₂tBu,
-CH₂NHCH₂CH₂SO₂CH₃,
-CH₂NH(CH₂CO₂H),
-CH₂N(C(O)Me)(CH₂CO₂H),
-CH₂-N-pyrrolidin-2-one,
-CH₂-(1-methylpyrrolidin-2-one-3-yl),
-CH₂S(O)Me,

-CH₂S(O)Et,
-CH₂S(O)₂Me,
-CH₂S(O)₂Et,
-CH₂S(O)iPr,
-CH₂S(O)₂iPr,
-CH₂S(O)tBu,
-CH₂S(O)₂tBu,
-CH₂CO₂H, CH₂C(O)NH₂,
-CH₂C(O)NMe₂,
-CH₂C(O)NHMe,
-CH₂C(O)-N-pyrrolidine,
-CH₂S(O)₂Me, CH₂S(O)Me,
-CH(OH) CO₂H,
-CH(OH)C(O)NH₂,
-CH(OH)C(O)NHMe,
-CH(OH)C(O)NMe₂,
-CH(OH)C(O)NEt₂,
-CH₂CH₂CO₂H,
-CH₂CH₂CO₂Me,
-CH₂CH₂CO₂Et,
-CH₂CH₂C(O)NH₂,
-CH₂CH₂C(O)NHMe,
-CH₂CH₂C(O)NMe₂,
-CH₂CH₂-5-tetrazolyl,
-CH₂CH₂S(O)₂Me,
-CH₂CH₂S(O)Me,
-CH₂CH₂S(O)₂Et,
-CH₂CH₂S(O) Et,
-CH₂CH₂S(O)iPr,

$-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{iPr},$
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{tBu},$
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{tBu},$
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{NH}_2,$
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{NHMe},$
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{NMe}_2,$
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{NH}_2,$
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{NHMe}$
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{NMe}_2,$
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{Me},$
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{Et},$
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{Me},$ or
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{Et}.$

The compound of formula IB having as preferred substituents;;

R and R' are independently methyl or ethyl;

RP, RB, RB₄, and RT₃ are independently, hydrogen or methyl;

RP₃ and RB₇ are independently hydrogen, methyl, ethyl, -O-methyl, or cyclopropyl;

(L_{P1}) and (L_{BT}) divalent linking groups are both bonds;

(L_{P2}) is a bond, -CH₂-, -CH(OH)-, or -C(Me)OH-;

Z_P is 1,1-dimethylethyl; 1-hydroxycyclopentyl, 1-hydroxycyclohexyl, 3-ethyl-3-hydroxypentyl, 3-ethyl-3-hydroxypentenyl, 3-ethyl-3-hydroxypentynyl;

Z_{BT} is

$-\text{CO}_2\text{H},$
 $-\text{CO}_2\text{Me},$
 $-\text{CO}_2\text{Et},$
 $-\text{C}(\text{O})\text{CH}_2\text{S}(\text{O})\text{Me},$
 $-\text{C}(\text{O})\text{CH}_2\text{S}(\text{O})\text{Et},$
 $-\text{C}(\text{O})\text{CH}_2\text{S}(\text{O})_2\text{Me},$
 $-\text{C}(\text{O})\text{CH}_2\text{S}(\text{O})_2\text{Et},$

-C(O)CH₂CH₂S(O)Me,
-C(O)CH₂CH₂S(O)Et,
-C(O)CH₂CH₂S(O)₂Me,
-C(O)CH₂CH₂S(O)₂Et,
-C(O)CH(Me)CH₂CO₂H,
-C(O)CH(Me)CH₂CO₂Me,
-C(O)CH(Me)CH₂CO₂Et,
-C(O)CH(Me)CH₂CO₂iPr,
-C(O)CH(Me)CH₂CO₂tBu,
-C(O)CH(Me)CH(Me)CO₂H,
-C(O)CH(Me)CH(Me)CO₂Me,
-C(O)CH(Me)CH(Me)CO₂Et,
-C(O)CH(Me)CH(Me)CO₂iPr,
-C(O)CH(Me)CH(Me)CO₂tBu,
-C(O)CH(Me)C(Me)₂CO₂H,
-C(O)CH(Me)C(Me)₂CO₂Me,
-C(O)CH(Me)C(Me)₂CO₂Et,
-C(O)CH(Me)C(Me)₂CO₂iPr,
-C(O)CH(Me)C(Me)₂CO₂tBu,
-C(O)CH(Me)CH(Et)CO₂H,
-C(O)CH(Me)CH(Et)CO₂Me,
-C(O)CH(Me)CH(Et)CO₂Et,
-C(O)CH(Me)CH(Et)CO₂iPr,
-C(O)CH(Me)CH(Et)CO₂tBu,
-C(O)C(O)OH,
-C(O)C(O)NH₂,
-C(O)C(O)NHMe,
-C(O)C(O)NMe₂,
-C(O)NH₂,

-C(O)NMe₂,
-C(O)NH-CH₂-C(O)OH,
-C(O)NH-CH₂-C(O)OMe,
-C(O)NH-CH₂-C(O)OEt,
-C(O)NH-CH₂-C(O)OiPr,
-C(O)NH-CH₂-C(O)OtBu,
-C(O)NH-CH(Me)-C(O)OH,
-C(O)NH-CH(Me)-C(O)OMe,
-C(O)NH-CH(Me)-C(O)OEt,
-C(O)NH-CH(Me)-C(O)iPr,
-C(O)NH-CH(Me)-C(O)tBu,
-C(O)NH-CH(Et)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OMe,
-C(O)NH-C(Me)₂-C(O)OEt,
-C(O)NH-C(Me)₂-C(O)iPr,
-C(O)NH-C(Me)₂-C(O)tBu,
-C(O)NH-CMe(Et)-C(O)OH,
-C(O)NH-CH(F)-C(O)OH,
-C(O)NH-CH(CF₃)-C(O)OH,
-C(O)NH-CH(OH)-C(O)OH,
-C(O)NH-CH(cyclopropyl)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-CF(Me)-C(O)OH,
-C(O)NH-C(Me)(CF₃)-C(O)OH,
-C(O)NH-C(Me)(OH)-C(O)OH,
-C(O)NH-C(Me)(cyclopropyl)CO₂H
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH₂-C(O)OMe,
-C(O)NMe-CH₂-C(O)OEt,
-C(O)NMe-CH₂-C(O)OiPr,
-C(O)NMe-CH₂-C(O)tBu,

-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH(Me)-C(O)OH,
-C(O)NMe-CH(F)-C(O)OH,
-C(O)NMe-CH(CF₃)-C(O)OH,
-C(O)NMe-CH(OH)-C(O)OH,
-C(O)NMe-CH(cyclopropyl)-C(O)OH,
-C(O)NMe-C(Me)₂-C(O)OH,
-C(O)NMe-CF(Me)-C(O)OH,
-C(O)NMe-C(Me)(CF₃)-C(O)OH,
-C(O)NMe-C(Me)(OH)-C(O)OH,
-C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,
-C(O)NHS(O)Me,
-C(O)NHSO₂Me,
-C(O)-NH-5-tetrazolyl,
-C(O)NHS(O)Me,
-C(O)NHS(O)Et,
-C(O)NHSO₂Me,
-C(O)NHSO₂Et,
-C(O)NHS(O)iPr,
-C(O)NHSO₂iPr,
-C(O)NHS(O)tBu,
-C(O)NHSO₂tBu,
-C(O)NHCH₂S(O)Me,
-C(O)NHCH₂S(O)Et,
-C(O)NHCH₂SO₂Me,
-C(O)NHCH₂SO₂Et,
-C(O)NHCH₂CH₂S(O)Me,
-C(O)NHCH₂CH₂S(O)Et,
-C(O)NHCH₂CH₂SO₂Me,
-C(O)NHCH₂CH₂SO₂Et,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)SO₂Me,


-C(O)-N(Me)-5-tetrazolyl,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)S(O)Et,
-C(O)N(Me)SO₂Me,
-C(O)N(Me)SO₂Et,
-C(O)N(Me)S(O)iPr,
-C(O)N(Me)SO₂iPr,
-C(O)N(Me)S(O)tBu,
-C(O)N(Me)SO₂tBu,
-C(O)N(Me)CH₂S(O)Me,
-C(O)N(Me)CH₂S(O)Et,
-C(O)N(Me)CH₂SO₂Me,
-C(O)N(Me)CH₂SO₂Et,
-C(O)N(Me)CH₂CH₂S(O)Me,
-C(O)N(Me)CH₂CH₂S(O)Et,
-C(O)N(Me)CH₂CH₂SO₂Me,
-C(O)N(Me)CH₂CH₂SO₂Et,
-CH₂CO₂H,
-CH₂-5-tetrazolyl,
-CH₂CO₂Me,
-CH₂CO₂Et,
-CH₂NHS(O)Me,
-CH₂NHS(O)Et,
-CH₂NHSO₂Me,
-CH₂NHSO₂Et,
-CH₂NHS(O)iPr,
-CH₂NHSO₂iPr,
-CH₂NHS(O)tBu,
-CH₂NHSO₂tBu,
-CH₂NHCH₂CH₂SO₂CH₃,

-CH₂NH(CH₂CO₂H),
-CH₂N(C(O)Me)(CH₂CO₂H),
-CH₂-N-pyrrolidin-2-one,
-CH₂-(1-methylpyrrolidin-2-one-3-yl),
-CH₂S(O)Me,
-CH₂S(O)Et,
-CH₂S(O)₂Me,
-CH₂S(O)₂Et,
-CH₂S(O)iPr,
-CH₂S(O)₂iPr,
-CH₂S(O)tBu,
-CH₂S(O)₂tBu,
-CH₂CO₂H, CH₂C(O)NH₂,
-CH₂C(O)NMe₂,
-CH₂C(O)NHMe,
-CH₂C(O)-N-pyrrolidine,
-CH₂S(O)₂Me, CH₂S(O)Me,
-CH(OH)CO₂H,
-CH(OH)C(O)NH₂,
-CH(OH)C(O)NHMe,
-CH(OH)C(O)NMe₂,
-CH(OH)C(O)NEt₂,
-CH₂CH₂CO₂H,
-CH₂CH₂CO₂Me,
-CH₂CH₂CO₂Et,
-CH₂CH₂C(O)NH₂,
-CH₂CH₂C(O)NHMe,
-CH₂CH₂C(O)NMe₂,
-CH₂CH₂-5-tetrazolyl,

-CH₂CH₂S(O)₂Me,
-CH₂CH₂S(O)Me,
-CH₂CH₂S(O)₂Et,
-CH₂CH₂S(O) Et,
-CH₂CH₂S(O)iPr,
-CH₂CH₂S(O)₂iPr,
-CH₂CH₂S(O)tBu,
-CH₂CH₂S(O)₂tBu,
-CH₂CH₂S(O)NH₂,
-CH₂CH₂S(O)NHMe,
-CH₂CH₂S(O)NMe₂,
-CH₂CH₂S(O)₂NH₂,
-CH₂CH₂S(O)₂NHMe
-CH₂CH₂S(O)₂NMe₂,
-CH₂CH₂CH₂S(O)Me,
-CH₂CH₂CH₂S(O)Et,
-CH₂CH₂CH₂S(O)₂Me, or
-CH₂CH₂CH₂S(O)₂Et.

Particularly preferred compounds of the invention and salts and prodrug derivatives are represented by formulae C1 to C22 as follows:


C1)




C2)

P-16541


70




C3)




C4)




C5)



C6)




C7)




C8)

P-16541


71




C9)




C10)



C11)




C12)




C13)

P-16541


72




C14)




C15)



C16)




C17)




C18)

P-16541


73




C19)




C20)




C21)



C22)



Particularly preferred is the compound represented by the structural formula AA or a pharmaceutically acceptable salt or prodrug thereof:



EXAMPLES

General Experimental Conditions:

The starting material/intermediate is the compound from the immediate preceding experimental unless otherwise indicated.

All reactions are performed under nitrogen/argon atmosphere, in a stirred reaction vessel, and at room temperature unless indicated otherwise.

Unless otherwise indicated, the organic layer is $\text{MgSO}_4/\text{Na}_2\text{SO}_4$ dried is defined as stirring the solution with a dessicant for 5-15 m and filtering off the dessicant to give an anhydrous filtrate.

For analogous multi-step reaction procedures, the yield is given either for the ultimate step or overall multi-steps as indicated.

Solutions are "concentrated" at a range of 25-75 °C with reduced pressure. in-vacuo – 25-75 °C; 0.05 to 1 mm

Unless otherwise indicated, "the residue is chromatographed" is defined as silica gel chromatography of residue with moderate nitrogen pressure (flash chromatography) or a medium pressure chromatography systems using a silica gel to crude product ratio of ~10-100.

For HPLC, the conditions listed are for the analytical trace only. For Preparative HPLC, the eluent is similar to analytical HPLC eluent.

Thin layer chromatography is performed with silica gel plates with UV and/or appropriate staining solution.

NMR spectra are obtained with either 300 or 400 mHz spectrometer.

NMR data is listed to denote spectrum is consistent with assigned structure.

"NMR" notation without data denotes spectrum is consistent with assigned structure.

HRMS – high resolution mass spectrum

ES-MS – electrospray mass spectrum

Abbreviations:

Aq - aqueous

d – day

eq - equivalent

h – hour

m – minute

satd – saturated

disp – dispersion

quant – quantitative

rt for retention time (both small caps to minimize confusion with RT)

RT – room temperature

Chemical Definitions:

BF₃-OEt₂ – boron trifluoride etherate

BnBr – benzyl bromide

CH₂Cl₂-dichloromethane

CH₃CN – acetonitrile

CO-carbon monoxide

CsF – cesium fluoride

DMAP - 4-(dimethylamino)pyridine

DMF – N,N-dimethylformamide

DMSO – dimethylsulfoxide

DPPB – 1,4-bis(diphenylphosphino)butane

DPPF – dichloro[1,1'-bis(diphenylphosphino)ferrocene

EDCI – 3-Ethyl-1-[3-(dimethylamino)propyl]carbodiimide

hydrochloride

Et₃N – triethylamine

EtMgBr- ethyl magnesium bromide

EtOAc – ethyl acetate

EtOH – ethanol

H₂ – hydrogen pressure

H₂NCH₂CO₂Me – methyl glycinate
Hept – heptane
Hex - hexanes
HN(OMe)Me – N-methyl-O-methyl hydroxylamine
HNMe₂ – dimethyl amine
HATU – O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate
HOAT – 7-aza-1-hydroxybenzotriazole
HOBT – 1-hydroxybenzotriazole
K₂CO₃ – potassium carbonate
KI – potassium iodine
KOH – potassium hydroxide
LAH – lithium aluminum hydride
LiHMDS – lithium hexamethyldisilazide
Lindlar catalyst – Pd-CaCO₃-PbO
LiOH – lithium hydroxide
mCPBA – meta-chloroperbenzoic acid
MeI – methyl iodide
MeOH – methanol
NaBH₄ – sodium borohydride
MgSO₄- magnesium sulfate
NaH – sodium hydride
NaHCO₃-sodium bicarbonate
NaI – sodium iodide
Na₂SO₄- sodium sulfate
NH₄Cl- ammonium chloride
NMO – 4-methylmorpholine N-oxide
NMP - N-methylpyrrolidin-2-one
Na-S-R₃ – sodium alkylmercaptide
PBr₃ – phosphorus tribromide
Pd(DPPF) - palladium dichloro[1,1'-bis(diphenylphosphino)ferrocene]
Pd(OAc)₂ – palladium (II) acetate
Pd(TPP)₄ – palladium tetrakis(triphenylphosphine)
Pd-C – palladium on carbon

Pd-C/H₂ – palladium on carbon with hydrogen pressure

pTSA – para-toluenesulfonic acid

Pyr – pyridine

Red-Al – sodium bis(2-methoxyethoxy)aluminum hydride

R₂MgBr – alkyl magnesium bromide

R₃MgBr – alkyl magnesium bromide

R₅MgBr – alkyl magnesium bromide

R₃S(O)₂Cl – alkylsulfonyl chloride

R₂S(O)₂NH₂ – alkylsulfonamide

TBSCl- *tert*-butyldimethylsilyl chloride

tBuC(O)CH₂Br – 1-bromopinacolone

Tf₂O – triflic anhydride

TFA – trifluoroacetic acid

THF – tetrahydrofuran


Ti(OiPr)₄ – titanium tetraisopropoxide

TMS-acetylene – trimethylsilyl acetylene


TPAP – tetrapropylammonium perruthenate

Zn(OTf)₂ – zinc trifluoromethane sulfonate


Scheme 1.




Scheme II.




Here, $R'-CH_2$ means R'
minus a methylene group




Scheme III.



Scheme IV.



Scheme V.



Scheme I. The known 2-fluoro-4-iodo-3-trimethylsilanyl-benzaldehyde is reacted with a mercaptoacetic acid ester to form benzo[b]thiophene ester 1. Benzo[b]thiophene 1. is reacted with an alkyl Grignard reagent (0.9 to 1.3 equivalents) at low temperature in diethylether or THF to undergo magnesium exchange with the iodine of 1. that is subsequently treated with a disubstituted ketone to produce the tert-carbinol 6. tert-Carbinol 6. is reacted with an ortho-substituted phenol in the presence of a Lewis acid, e.g., borontrifluoride etherate from 0 °C. to room temperature to give the diarylmethane 7. The free hydroxyl of diarylmethane 7. is alkylated with an alpha-halo ketone in a polar aprotic solvent such as acetone or acetonitrile, with or without catalytic potassium iodide from room temperature to the reflux temperature of the mixture. The alkylated intermediate preceding is saponified with lithium, potassium, or sodium hydroxide and alkanol from room temperature to the reflux temperature of the mixture to produce the acid 8. The ketone fragment of acid 8. is reduced with lithium, potassium, or sodium borohydride or cyanoborohydride (1 to 4 equivalents) in alkanol to the carbinol 9.

Alternately, benzo[b]thiophene 1. can be used as follows: 1. is reacted with an excess of alkyl Grignard reagent (3-6 equivalents) in THF or ether at higher temperature, e.g., room temperature to not only undergo magnesium exchange with the iodine of 1. but also to produce the tert-carbinol fragment of 2. The mixture is subsequently treated with excess carbon dioxide to produce the acid 2. Reaction of acid 2. with an ortho-substituted phenol in the presence of a Lewis acid (0.01 to 5 equivalents), e.g., boron trifluoride etherate from 0 °C. to room temperature gives a diarylmethane acid 3 that is esterified with acid and alkanol to the ester 3. The free hydroxyl of ester 3. is alkylated with an alpha-halo ketone in polar aprotic solvent to give an intermediate which is subsequently saponified with lithium, potassium, or sodium hydroxide in an alkanol (room temperature to reflux temperature) to give the acid 4. Acid 4. is reduced to the carbinol 5. with lithium, potassium, or sodium borohydride or cyanoborohydride (1 to 4 equivalents) in alkanol (0 ° C. to room temperature).

Scheme II. Methyl, p-fluorobenzoate is reacted with excess alkyl Grignard reagent (2 to 5 equivalents) to produce the tert-carinol intermediate, which eliminates water upon workup to give the olefin 10. Ortho metallation with an alkyllithium reagent of olefin 10. in diethylether or THF from 0 °C. to room temperature followed by treatment with DMF produces the aldehyde 11. Aldehyde 11. is reacted with a mercaptoacetic acid ester to give the benzo[b]thiophene 12. The benzo[b]thiophene 12. will alkylate an ortho-substituted phenol in the presence of a Lewis acid, e.g., boron trifluoride etherate (0.01 to 5 equivalents) to give the diarylmethane 13. The free hydroxyl of diarylmethane 13. is alkylated with an alpha-halo ketone in a polar aprotic solvent, e.g., acetone or acetonitrile from room temperature to the reflux temperature of the mixture, and the alkylated intermediate is subsequently saponified with lithium, potassium, or sodium hydroxide in an alkanol (room temperature to the reflux temperature of the mixture) to give the acid 14. The acid 14. is reduced to the carbinol 15. with lithium, potassium, or sodium borohydride or cyanoborohydride (1-4 equivalents) in alkanol from 0 ° C. to room temperature.

Scheme III. Benzo[b]thiophene 16. is prepared by the reaction of a mercaptoacetic acid ester and 2-fluoro-5-methoxy-benzaldehyde in DMF. The

benzo[b]thiophene 16. is reacted with excess alkyl Grignard reagent (2 to 5 equivalents) in diethylether or THF (0 ° C. to room temperature) to give the tert-carbinol 17. The tert-carbinol 17. alkylates an ortho-substituted phenol in the presence of a Lewis acid, e.g., boron trifluoride etherate (0.01 to 5 equivalents) to give the diarylmethane 18. The free hydroxyl of diarylmethane 18. is alkylated with an alpha-halo ketone or ester in a polar aprotic solvent, e.g., acetone or acetonitrile from room temperature to the reflux temperature of the mixture to give the methoxy intermediate 19. The intermediate 19. is demethylated with boron tribromide in halocarbon solvent from 0 ° to room temperature to give the free hydroxyl intermediate 20. The intermediate 20. is converted to the triflate 21. with triflic anhydride and base, and the triflate 21. is reacted with carbon monoxide (1-100 psi) in the presence of palladium catalysis (0.01 to 0.10 equivalents) , e.g., palladium acetate and DPPB, and base to give the ester intermediate which is subsequently saponified with lithium, potassium, or sodium hydroxide in alkanol to give the keto acid 22. The keto acid 22. is reduced to the carbinol 23. with lithium, potassium, or sodium borohydride or cyanoborohydride (1-4 equivalents) in alkanol from 0 ° C. to room temperature.

Scheme IV. Intermediate 19 is reacted with an alkyl Grignard reagent (1-5 equivalents) in ether or THF to give the methoxy carbinol 24. Methoxy carbinol 24. is demethylated with sodium mercaptoethylate in a polar aprotic solvent, e.g., DMF from room temperature to 150 ° C. to give the hydroxy carbinol 25. The hydroxy carbinol 25. is converted to the triflate 26. with triflic anhydride and base, and the triflate 26. is reacted with carbon monoxide (1-100 psi) in the presence of palladium catalysis (0.01 to 0.10 equivalents) , e.g., palladium acetate and DPPB, and base to give the ester 27. The ester 27. is saponified with lithium, potassium, or sodium hydroxide in alkanol to give the acid 28.


Scheme V. The free hydroxyls of esters 3., 7., 13., and 18., from Schemes I. II. And III. above are converted to the triflates 29. with triflic anhydride and base. Each of these triflates 29. is couple to TMS-acetylene in the presence of a palladium catalyst, e.g., PdCl₂(PPh₃)₂ (0.01 to 10%) in a polar aprotic solvent, e.g., acetonitrile from room temperature to 150 C. to give the TMS protected acetylenes as intermediates, that are deprotected with fluoride ion in THF to give acetylenes 30.

Each of the acetylenes 30. are deprotonated with, e.g., LiHMDS and treated with a ketone to produce the carbinols 31. Each of the carbinols 31. are saponified with lithium, potassium, or sodium hydroxide in alkanol to give the acids 32.


Each of the free acids produced in each of the above schemes (3., 4., 5., 8., 9., 14., 15., 22., 23., and 28.) are converted to lower alkanol esters and lower alkyl secondary or tertiary carboxamides using reactions well known to those skilled in the art.

Example 1.

Preparation of 6-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid.




A. 6-Iodo-benzo[b]thiophene-2-carboxylic acid ethyl ester



A solution of 2-fluoro-4-iodo-3-trimethylsilylanyl-benzaldehyde (14.2 g, 44.1 mmol) (*Tetrahedron Letters* 1992, p7499-7502) in DMF (50 mL) is treated with mercapto-acetic acid ethyl ester (5.80 mL, 52.9 mmol) and K_2CO_3 (12.2 g, 88.2 mmol). The resulting suspension is stirred at 75 °C for 60 min and quenched with water (300 mL). The mixture is extracted with EtOAc (3 x 50 mL,) and the organic layer is dried over Na_2SO_4 , concentrated, and purified by silica gel column chromatography (5-20% EtOAc/Hex), to give the title compound as an oil (9.40 g, 64%).

¹H-NMR (ppm, CDCl₃), δ: 8.21 (1 H, d, *J*=1.8 Hz), 7.98 (1 H, s), 7.67 (1 H, dd, *J*=1.8, 8.8 Hz), 7.58 (1 H, d, *J*=8.4 Hz), 4.40 (2 H, q, *J*=7.0 Hz), 1.42 (3 H, t, *J*=7.0 Hz).


B. 6-(1-Ethyl-1-hydroxy-propyl)-benzo[b]thiophene-2-carboxylic acid ethyl ester.



To a stirred solution of 6-iodo-benzo[b]thiophene-2-carboxylic acid ethyl ester (2.95 g, 8.88 mmol) in THF (60 mL) at -78°C is added ethylmagnesium bromide (4.45 mL, 3.0 M). The reaction mixture is stirred for 15 min, and 3-pentanone (3.0 mL) is added. The mixture is allowed to warm to RT and stirred for 30 min. The reaction is quenched with HCl (15 mL, 1.0 M) and extracted with EtOAc (2 x 50 mL). The organic layer is dried over Na_2SO_4 , concentrated, and purified by silica gel column chromatography to give the title compound (0.65 g, 25%).

H-NMR (ppm, CDCl_3), d: 8.02 (1 H, s), 7.95 (1 H, s), 7.80 (1 H, d, $J=8.8$ Hz), 7.35 (1 H, dd, $J=1.3, 8.4$ Hz), 4.40 (2 H, q, $J=7.0$ Hz), 1.88 (4 H, q, $J=7.5$ Hz), 1.42 (3 H, t, $J=7.0$ Hz), 0.78 (6 H, q, $J=7.5$ Hz).


C. 6-[1-Ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-benzo[b]thiophene-2-carboxylic acid ethyl ester.



To a mixture of 6-(1-ethyl-1-hydroxy-propyl)-benzo[b]thiophene-2-carboxylic acid ethyl ester (0.650 g, 2.23 mmol) and o-cresol (0.480 g, 4.45 mmol) in CH_2Cl_2 (20 mL) at -78°C is added $\text{BF}_3\cdot\text{Et}_2\text{O}$ (0.316 g, 2.23 mol). After stirring for 10 min, the reaction mixture is allowed to warm to 0°C over 30 min. The reaction is quenched with water (15 mL) and extracted with EtOAc (2 x 50 mL). The organic layer is dried over Na_2SO_4 , concentrated, and purified by silica gel column chromatography to give the title compound (0.780 g, 92%).

MS (ES) m/e 383.1 ($\text{M}+1$), 381.1 ($\text{M}-1$).

D. 6-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid ethyl ester.



A solution of 6-[1-ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-benzo[b]thiophene-2-carboxylic acid ethyl ester (0.780 g, 2.04 mmol) and 1-bromopinacolone (0.731 g, 4.08 mmol) in acetone (30 mL) is treated with K_2CO_3

(0.563 g, 4.08 mmol) and stirred at RT for 4 h. The mixture is filtered, and the filtrate is concentrated. The residue is purified using silica gel column chromatography (10-15% EtOAc/Hex) to provide the title compound (0.89 g, 91%).

MS (ES) *m/e* 498.2 (M+18).


E. 6-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid.

A solution of 6-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid ethyl ester (0.89 g, 1.86 mmol) in MeOH (5.0 mL) and THF (5.0 mL) and is treated with NaOH (2.0 M, 10.0 mL). The resulting mixture is stirred at RT for 2 h. The mixture is concentrated, acidified with HCl (1 N) until pH~3, and extracted with EtOAc (2 x 50 mL). The organic layer is dried over Na₂SO₄, concentrated, and purified using silica gel column chromatography (50% EtOAc/Hex) to afford the title compound (840 mg, 99%).

MS (ES) *m/e* 451.1(M-1), 470.2 (M+18).

Example 2.

Preparation of 6-{1-ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-2-carboxylic acid.




A solution of 6-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid (102 mg, 0.226 mmol) in THF (5.0 mL) at RT is treated with NaBH₄ (17 mg, 0.451 mmol). The resulting mixture is stirred for 4 h. The reaction is quenched with HCl (1.0 N, 1.0 mL) and extracted with EtOAc (2 x 20 mL). The organic layer is dried over Na₂SO₄ and concentrated to afford the title compound (90 mg, 88%).

MS (ES) *m/e* 453.1(M-1), 472.2 (M+18).

Example 3.


Preparation of 6-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid dimethylamide.



6-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid (290 mg, 0.642 mmol) in CH_2Cl_2 (5.0 mL) is treated with DMAP (235 mg, 1.93 mmol) and EDC (184 mg, 0.962 mmol). The mixture is stirred for 15 min at RT, and dimethylamine hydrochloride (78 mg, 0.962 mmol) is added. The reaction is stirred for 18 h and quenched with aqueous NH_4Cl (5.0 mL). The organic layer is loaded onto a silica gel column and purified with 50% EtOAc/Hex to afford the title compound (280 mg, 91%).
MS (ES) m/e 480.2 ($M+1$).

Example 4.


Preparation of 6-{1-ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-2-carboxylic acid dimethylamide.



6-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid dimethylamide (235 mg, 0.490 mmol) is reduced by NaBH_4 (37 mg, 0.980 mmol) as in Step F11 to afford the title compound (230 mg, 100%).
MS (ES) m/e 482.2 ($M+1$).

Example 5.

Preparation of [(6-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carbonyl)-amino]-acetic acid.



6-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid (270 mg, 0.596 mmol) in CH_2Cl_2 (5.0 mL) is


treated with DMAP (218 mg, 1.79 mmol) and EDC (172 mg, 0.895 mmol). The mixture is stirred for 15 min at RT, and methyl ester hydrochloride (112 mg, 0.895 mmol) is added. The reaction is stirred for 18 h and quenched with NH_4Cl (5.0 mL). The organic layer is loaded onto a silica gel column and purified with 20-50% EtOAc/Hex to afford the intermediate amide ester.

The intermediate is dissolved in methanol (3.0 mL) and THF (2.0 mL) and treated with NaOH (2.0 M, 5.0 mL). The resulting mixture is stirred at RT for 3 h. The mixture is concentrated, acidified with HCl (1 N) to pH~3, and extracted with EtOAc (2 x 20 mL). The organic layer is dried over Na_2SO_4 and concentrated to afford the title compound (285 mg, 94%).

MS (ES) m/e 510.1 (M+1), 508.1 (M-1).

Example 6.

Preparation of [(6-{1-ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-2-carbonyl)-amino]-acetic acid.




[(6-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carbonyl)-amino]-acetic acid (220 mg, 0.432 mmol) is reduced by NaBH_4 (33 mg, 0.863 mmol) as in Step F11 to afford the title compound (175 mg, 79%).


MS (ES) m/e 512.1 (M+1), 510.1 (M-1).

Example 7.

Preparation of 2-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carboxylic acid.




A. 2-(1-Ethyl-1-hydroxy-propyl)-benzo[b]thiophene-6-carboxylic acid.



A solution of 6-iodo-benzo[b]thiophene-2-carboxylic acid ethyl ester (5.05 g, 15.2 mmol) in THF (200 mL) at -78°C is de-oxygenated (vacuum/ N_2 purge 3 x). Ethylmagnesium bromide (25.3 mL, 3.0 M) is added, and the reaction mixture is allowed to warm to RT over 30 min. The mixture is stirred for 10 min at RT, cooled to -78°C , and treated with clean dry ice (100 g). The mixture is allowed to warm to RT over 60 min and is quenched with HCl (50 mL, 1.0 M). THF is removed under vacuum, and the residue is extracted with EtOAc (3 x 100 mL). The organic layer is dried over Na_2SO_4 and concentrated to give the product (3.80 g, 94%).

MS (ES) m/e 263.0 (M-1).


B. 2-[1-Ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-benzo[b]thiophene-6-carboxylic acid.



A solution of 2-(1-ethyl-1-hydroxy-propyl)-benzo[b]thiophene-6-carboxylic acid (3.80 g, 14.4 mmol) and o-cresol (1.94 g, 18.0 mmol) in CH_2Cl_2 (100 mL) at -40°C is added $\text{BF}_3\cdot\text{Et}_2\text{O}$ (1.02 g, 7.20 mol). After stirring for 10 min, the reaction mixture is allowed to warm up to 0°C over 30 min. The reaction is quenched with water (15 mL) and the aqueous layer is extracted with EtOAc (50 mL). The organic layer is dried over Na_2SO_4 , concentrated, and purified by silica gel column chromatography (2% HOAc in 50% EtOAc/Hex) to give the product (3.98 g, 78%).


MS (ES) m/e 353.0 (M-1).

C. 2-[1-Ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-benzo[b]thiophene-6-carboxylic acid methyl ester.



A solution of 2-[1-ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-benzo[b]thiophene-6-carboxylic acid (3.98 g, 11.23 mmol) in MeOH (100 mL) is treated with H_2SO_4 (concentrated, 1.0 mL). The mixture is stirred at 80 °C for 8 h and neutralized with aqueous NaHCO_3 . The MeOH is removed under vacuum and the residue is extracted with EtOAc (2 x 100 mL). The combined organic layer is dried over Na_2SO_4 , concentrated and purified by silica gel column chromatography (20% EtOAc/Hex) to give the product (4.05 g, 98%). MS (ES) m/e 367.1 (M-1).

D. 2-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carboxylic acid methyl ester.




A solution of 2-[1-ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-benzo[b]thiophene-6-carboxylic acid methyl ester (4.05 g, 11.0 mmol) and 1-bromopinacolone (2.36 g, 13.2 mmol) in acetone (100 mL) is treated with K_2CO_3 (3.04 g, 22.0 mmol) and stirred at RT for 16 h. The mixture is filtered, and the filtrate is concentrated. The residue is purified by silica gel column chromatography (10-15% EtOAc/Hex) to provide the title compound (4.50 g, 88%). MS (ES) m/e 484.2 (M+18).

E. 2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carboxylic acid.

2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carboxylic acid methyl ester (120 mg, 0.257 mmol) is hydrolyzed as in Step E11 to give the title compound (72 mg, 62%). MS (ES) m/e 451.2 (M-1), 470.2 (M+18).

Example 8.

Preparation of 2-{1-Ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-6-carboxylic acid.




2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carboxylic acid (53 mg, 0.117 mmol) is reduced by NaBH_4 (9 mg, 0.234 mmol) as in a reaction analogous to Example 2 to afford the title compound (53 mg, 99%).

MS (ES) m/e 453.2 (M-1), 472.2 (M+18).

Example 9.

Preparation of 2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carboxylic acid dimethylamide.




2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carboxylic acid (175 mg, 0.387 mmol), DMAP (142 mg, 1.16 mmol), EDC (111 mg, 0.580 mmol), and dimethylamine hydrochloride (47 mg, 0.580 mmol) are reacted analogous to Example 3 to afford the title compound (140 mg, 76%).

MS (ES) m/e 480.2 (M+1).

Example 10.

Preparation of 2-{1-Ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-6-carboxylic acid dimethylamide.




2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carboxylic acid dimethylamide (90 mg, 0.188 mmol) is reduced using NaBH_4 (14 mg, 0.376 mmol) in a reaction analogous to Example 2 to afford the title compound (90 mg, 100%).

MS (ES) m/e 482.2 (M+1).

Example 11.

Preparation of [(2-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carbonyl)-amino]-acetic acid.




2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carboxylic acid (175 mg, 0.387 mmol), DMAP (142 mg, 1.16 mmol), EDC (111 mg, 0.580 mmol), and glycine methyl ester hydrochloride (73 mg, 0.580 mmol) are reacted and hydrolyzed in a sequence analogous to Example 5 to afford the title compound (180 mg, 91%).

MS (ES) m/e 508.2 (M-1), 510.1 (M+1).

Example 12.

Preparation of [(2-{1-Ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-6-carbonyl)-amino]-acetic acid.




[(2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-6-carbonyl)-amino]-acetic acid (135 mg, 0.265 mmol) is reduced using NaBH_4 (20 mg, 0.530 mmol) as in example 2 to afford the title compound (135 mg, 99%).


MS (ES) m/e 512.2 (M+1), 510.2 (M-1).

Example 13.

Preparation of 5-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid.




A. 1-(1-Ethyl-propenyl)-4-fluoro-benzene.



A solution of 4-fluoro-benzoic acid methyl ester (19.1 g, 114 mmol) in THF (300 mL) at -78°C is treated with ethylmagnesium bromide (135 mL, 3.0 M). The reaction mixture is allowed to warm to 0°C over 30 min and quenched with HCl (350 mL, 1.0 M). THF is removed under vacuum, and the residue is extracted with EtOAc (3 x 100 mL). The organic layer is dried over Na_2SO_4 and concentrated. The intermediate tertiary alcohol is dissolved in CH_2Cl_2 (200 mL), cooled to -40°C , and treated with $\text{BF}_3\cdot\text{Et}_2\text{O}$ (11.5 g, 91.0 mol). The reaction mixture is allowed to warm to 0°C over 30 min and quenched with water (50 mL). The organic layer is dried over Na_2SO_4 and concentrated. The residue is purified using silica gel column chromatography (5% EtOAc/Hex) to afford the title compound (16.8 g, 90%).

H-NMR (ppm, CDCl_3), δ : 7.27 (2 H, m), 6.97 (2 H, m), 5.66 (1 H, q, $J=7.0$ Hz), 2.49 (2 H, q, $J=7.0$ Hz), 1.78 (3 H, d, $J=7.0$ Hz), 0.98 (3 H, t, $J=7.0$ Hz).

B. 5-(1-Ethyl-propenyl)-2-fluoro-benzaldehyde.




A solution of 1-(1-ethyl-propenyl)-4-fluoro-benzene (15.3 g, 93.2 mmol) in THF (150 mL) at -78°C is treated with *n*-BuLi (70 mL, 1.6 M). The reaction

mixture is allowed to warm to 0 °C over 60 min, cooled to -78 °C, and treated with DMF (10.2 g, 140 mmol). The mixture is stirred for 10 min and quenched with HOAc (10 mL) and water (200 mL). The mixture is allowed to warm to RT, and the THF is removed under vacuum. The residue is extracted with EtOAc (2 x 100 mL), and the organic layer is dried over Na₂SO₄ and concentrated. The crude product is purified using silica gel column chromatography (7% EtOAc/Hex) to afford the title compound (8.70 g, 49%).

H-NMR (ppm, CDCl₃), δ : 10.34 (1 H, s), 7.79 (1 H, dd, J =2.7, 6.6 Hz), 7.54 (1 H, m), 7.09 (1 H, dd, J =1.3, 9.7 Hz), 5.72 (1 H, q, J =7.0 Hz), 2.50 (2 H, q, J =7.5 Hz), 1.79 (3 H, d, J =7.0 Hz), 0.97 (3 H, t, J =7.5 Hz).


C. 5-(1-Ethyl-propenyl)-benzo[b]thiophene-2-carboxylic acid ethyl ester.



A solution of 5-(1-ethyl-propenyl)-2-fluoro-benzaldehyde (8.70 g, 45.3 mmol) in DMF (40 mL) is treated with mercapto-acetic acid ethyl ester (8.16 g, 67.9 mmol) and K₂CO₃ (12.5 g, 90.6 mmol). The resulting suspension is stirred at 80 °C for 60 min and quenched with water (300 mL). The mixture is extracted with EtOAc (2 x 200 mL), and the organic layer is dried over Na₂SO₄, concentrated, and purified using silica gel column chromatography (6% EtOAc/Hex), to give the title compound as an oil (11.0 g, 89%).


H-NMR (ppm, CDCl₃), δ : 8.01 (1 H, s), 7.78 (1 H, d, J =1.8 Hz), 7.75 (1 H, d, J =8.4 Hz), 7.44 (1 H, dd, J =1.8, 8.4 Hz), 5.75 (1 H, q, J =7.0 Hz), 4.41 (2 H, q, J =7.0 Hz), 2.58 (2 H, q, J =7.5 Hz), 1.82 (3 H, d, J =7.0 Hz), 1.42 (3 H, t, J =7.0 Hz), 0.97 (3 H, t, J =7.5 Hz).

D. 5-[1-Ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-benzo[b]thiophene-2-carboxylic acid ethyl ester.



5-(1-Ethyl-propenyl)-benzo[b]thiophene-2-carboxylic acid ethyl ester (5.50 g, 20.0 mmol), o-cresol (4.33 g, 40.0 mmol), and $\text{BF}_3\text{-Et}_2\text{O}$ (10 mL) are reacted analogous to Example 1C to afford the title compound (5.97 g, 78%). MS (ES) m/e 381.3 (M-1), 383.4 (M+1).

E. 5-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid ethyl ester.




5-[1-Ethyl-1-(4-hydroxy-3-methyl-phenyl)-propyl]-benzo[b]thiophene-2-carboxylic acid ethyl ester (5.97 g, 15.6 mmol), 1-bromopinacolone (4.20 g, 23.4 mmol), and K_2CO_3 (6.46 g, 46.8 mmol) in acetone (100 mL) are reacted analogous to Example 1D to provide the title compound (7.10 g, 95%). MS (ES) m/e 498.4 (M+18).

F. 5-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid.

5-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid ethyl ester (2.50 g, 5.20 mmol) is hydrolyzed as in Step E11 to give the title compound (2.20 g, 94%). MS (ES) m/e 451.2 (M-1), 470.2 (M+18).

Example 14.

Preparation of 5-{1-ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-2-carboxylic acid.




5-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid (0.280 g, 0.619 mmol) is reduced using NaBH_4 (14 mg, 0.376 mmol) analogous to Example 2 to afford the title compound (0.241 g, 86%).

MS (ES) m/e 453.1 (M-1), 472.2 (M+18).

Example 15.

Preparation of [(5-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carbonyl)-amino]-acetic acid.




5-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid (310 mg, 0.686 mmol), DMAP (167 mg, 1.37 mmol), EDC (197 mg, 1.03 mmol), and glycine methyl ester hydrochloride (172 mg, 1.37 mmol) are reacted and hydrolyzed in a sequence analogous to Example 5 to afford the title compound (150 mg, 43%).

MS (ES) m/e 508.2 (M-1), 510.1 (M+1).

Example 16.

Preparation of [(5-{1-ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-2-carbonyl)-amino]-acetic acid.




[(5-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carbonyl)-amino]-acetic acid (90 mg, 0.177 mmol) is reduced using NaBH_4 (13 mg, 0.354 mmol) in a reaction analogous to Example 2 to afford the title compound (73 mg, 81%).


MS (ES) m/e 510.2 (M-1), 512.2 (M+1).

Example 17.

Preparation of 2-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-5-carboxylic acid.




A. 5-Methoxy-benzo[b]thiophene-2-carboxylic acid ethyl ester.



A solution of 2-fluoro-5-methoxy-benzaldehyde (4.35 g, 28.2 mmol) in DMF (30 mL) is added mercapto-acetic acid ethyl ester (3.71 mL, 33.9 mmol) and K_2CO_3 (7.86 g, 57.0 mmol). The resulting suspension is stirred at 80 °C for 60 min and quenched with water (300 mL). The mixture is extracted with EtOAc (2 x 200 mL), and the organic layer is dried over Na_2SO_4 , concentrated, and purified by silica gel column chromatography (10% EtOAc/Hex), to give the title compound as an oil (4.40 g, 66%).

H-NMR (ppm, $CDCl_3$), δ : 7.97 (1 H, s), 7.70 (1 H, d, $J=8.8$ Hz), 7.27 (1 H, d, $J=2.6$ Hz), 7.09 (1 H, dd, $J=2.6, 8.8$ Hz), 4.41 (2 H, q, $J=7.0$ Hz), 3.88 (3 H, s), 1.42 (3 H, t, $J=7.0$ Hz).


B. 3-(5-Methoxy-benzo[b]thiophen-2-yl)-pentan-3-ol.



5-Methoxy-benzo[b]thiophene-2-carboxylic acid ethyl ester (4.13 g, 17.5 mmol) in THF (100 mL) at -78 °C is treated with ethylmagnesium bromide (17.5 mL, 3.0 M). The reaction mixture is allowed to warm to RT over 30 min and is stirred for 10 min. The mixture is quenched with water (40 mL) and acidified with HCl (50 mL, 1.0 M). THF is removed under vacuum, and the residue is extracted with EtOAc (2 x 40 mL). The organic layer is dried over Na_2SO_4 , concentrated, and purified using silica gel column chromatography (10% EtOAc/Hex) to give the title compound (4.53 g, 97%).

H-NMR (ppm, $CDCl_3$), δ : 7.63 (1 H, d, $J=8.8$ Hz), 7.17 (1 H, d, $J=2.2$ Hz), 7.03 (1 H, s), 6.92 (1 H, dd, $J=2.6, 8.8$ Hz), 3.86 (3 H, s), 1.91 (4 H, q, $J=7.0$ Hz), 0.91 (6 H, t, $J=7.0$ Hz).


C. 4-[1-Ethyl-1-(5-methoxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenol.



3-(5-Methoxy-benzo[b]thiophen-2-yl)-pentan-3-ol (2.16 g, 8.63 mmol), o-cresol (4.66 g, 43.1 mmol), and $\text{BF}_3\text{-Et}_2\text{O}$ (1.64 mL, 12.9 mmol) are reacted analogous to Example 1C to afford the title compound (2.29 g, 78%).

MS (ES) m/e 339.2 (M-1), 341.3 (M+1).


D. 1-{4-[1-Ethyl-1-(5-methoxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenoxy}-3,3-dimethyl-butan-2-one (2131468) (PF1-A03098-163)



4-[1-Ethyl-1-(5-methoxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenol (7.80 g, 22.9 mmol), 1-bromopinacolone (8.20 g, 45.8 mmol), and K_2CO_3 (6.32 g, 45.8 mmol) in acetone (100 mL) are reacted analogous to Example 1D to provide the title compound (9.20 g, 92%).

H-NMR (ppm, CDCl_3), δ : 7.53 (1 H, d, $J=8.8$ Hz), 7.15 (1 H, d, $J=2.6$ Hz), 7.02-7.06 (3 H, m), 6.87 (1 H, dd, $J=2.6, 8.8$ Hz), 6.50 (1 H, d, $J=8.4$ Hz), 4.84 (2 H, s), 3.85 (3 H, s), 2.25 (3 H, s), 2.15 (4 H, q, $J=7.5$ Hz), 1.26 (9 H, s), 0.73 (6 H, t, $J=7.5$ Hz).

E. 1-{4-[1-Ethyl-1-(5-hydroxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenoxy}-3,3-dimethyl-butan-2-one.




A solution of 1-{4-[1-ethyl-1-(5-methoxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenoxy}-3,3-dimethyl-butan-2-one (0.86 g, 1.96 mmol) in CH_2Cl_2 (15.0 mL) at 0 °C is treated with BBR_3 (3.0 mL, 2.94 mmol). The mixture is stirred for 2 h and quenched with water (10 mL). The aqueous layer is extracted with CH_2Cl_2 (30 mL). The organic layer is concentrated and purified using silica gel column chromatography (25% EtOAc/Hex) to afford the title compound (0.412 g, 50%).

H-NMR (ppm, CDCl_3), δ : 7.50 (1 H, d, $J=8.4$ Hz), 7.10 (1 H, d, $J=2.6$ Hz), 7.02-7.06 (2 H, m), 6.98 (1 H, s), 6.78 (1 H, dd, $J=2.6, 8.8$ Hz), 6.51 (1 H, d, $J=8.4$ Hz),

4.84 (2 H, s), 2.25 (3 H, s), 2.15 (4 H, q, $J=7.5$ Hz), 1.26 (9 H, s), 0.73 (6 H, t, $J=7.5$ Hz).


F. Trifluoromethanesulfonic acid 2-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophen-5-yl ester.



A solution of 1-{4-[1-ethyl-1-(5-hydroxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenoxy}-3,3-dimethyl-butan-2-one (3.21 g, 7.56 mmol) in CH_2Cl_2 (50 mL) at -78°C is treated with 2,6-lutidine (1.32 mL, 11.3 mmol) and trifluoromethanesulfonic acid anhydride (1.78 mL, 10.6 mmol). The mixture is stirred and allowed to warm to -20°C over 60 min. The mixture is quenched with water (5.0 mL). The organic layer is concentrated and purified using silica gel column chromatography (10% EtOAc/Hex) to afford the title compound (4.20 g, 99%).

$^1\text{H-NMR}$ (ppm, CDCl_3), δ : 7.69 (1 H, d, $J=8.8$ Hz), 7.57 (1 H, d, $J=2.2$ Hz), 7.13 (1 H, s), 7.11 (1 H, dd, $J=2.2, 8.8$ Hz), 7.04 (1 H, s), 7.01 (1 H, dd, $J=2.2, 8.8$ Hz), 6.52 (1 H, d, $J=8.4$ Hz), 4.84 (2 H, s), 2.27 (3 H, s), 2.15 (4 H, q, $J=7.5$ Hz), 1.26 (9 H, s), 0.74 (6 H, t, $J=7.5$ Hz).

G. 2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-5-carboxylic acid methyl ester.



A solution of trifluoromethanesulfonic acid 2-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophen-5-yl ester (4.10 g, 7.36 mmol) in DMSO (15 mL) and MeOH (10 mL) is treated with $\text{Pd}(\text{OAc})_2$ (0.169 g), DPPB (0.368 g) and Et_3N (5.3 mL). The mixture is stirred under CO (100 psi) at 80°C for 4 h. The MeOH is removed in vacuum, and the residue is poured into water (150 mL) and extracted with EtOAc (2 x 50 mL). The organic

layer is dried over Na_2SO_4 , concentrated, purified using silica gel column chromatography (10% EtOAc/Hex), to give the title compound as an oil (3.10 g, 90%).

$^1\text{H-NMR}$ (ppm, CDCl_3), δ : 8.38 (1 H, d, $J=2.6$ Hz), 7.87 (1 H, dd, $J=1.3, 8.4$ Hz), 7.70 (1 H, d, $J=8.8$ Hz), 7.19 (1 H, s), 7.05 (1 H, s), 7.01 (1 H, dd, $J=2.2, 8.4$ Hz), 6.52 (1 H, d, $J=8.4$ Hz), 4.84 (2 H, s), 3.94 (3 H, s), 2.26 (3 H, s), 2.18 (4 H, q, $J=7.5$ Hz), 1.26 (9 H, s), 0.74 (6 H, t, $J=7.5$ Hz).


MS (ES) m/e 484.4 ($\text{M}+18$).

H. 2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-5-carboxylic acid.

2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-5-carboxylic acid methyl ester (2.95 g, 6.32 mmol) is hydrolyzed as in Step E11 to give the title compound (2.80 g, 98%).
MS (ES) m/e 451.2 ($\text{M}-1$), 470.2 ($\text{M}+18$).

Example 18.


Preparation of 2-{1-ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-5-carboxylic acid.



2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-5-carboxylic acid (330 mg, 0.730 mmol) is reduced using NaBH_4 (55 mg, 1.46 mmol) in a reaction analogous to Example 2 to afford the title compound (0.306 g, 92%).
MS (ES) m/e 453.2 ($\text{M}-1$), 472.2 ($\text{M}+18$).

Example 19.


Preparation of [(2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-5-carbonyl)-amino]-acetic acid.



2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-5-carboxylic acid (310 mg, 0.686 mmol), DMAP (167 mg, 1.37 mmol), EDC (197 mg, 1.03 mmol), and glycine methyl ester hydrochloride (172 mg, 1.37 mmol) are reacted and hydrolyzed in a sequence analogous to Example 5 to afford the title compound (246 mg, 70%).
MS (ES) *m/e* 508.2 (M-1), 510.1 (M+1).

Example 20.


Preparation of [(2-{1-Ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-5-carbonyl)-amino]-acetic acid.



[(2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-5-carbonyl)-amino]-acetic acid (150 mg, 0.295 mmol) is reduced using NaBH₄ (22 mg, 0.589 mmol) analogous to Example 2 to afford the title compound (130 mg, 86%).
MS (ES) *m/e* 510.2 (M-1), 512.2 (M+1).

Example 21.

Preparation of 2-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-5-carboxylic acid dimethylamide.




A solution of 2-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-5-carboxylic acid (210 mg, 0.464 mmol) in CH₂Cl₂ (5.0 mL) is treated with Et₃N (1.0 mL), HOBT (94 mg, 0.70 mmol), and

EDC (134 mg, 0.70 mmol). The mixture is stirred for 10 min at RT, and dimethylamine (1.0 mL, 2.0 M in THF) is added. The reaction mixture is stirred for 18 h, loaded onto a silica gel column, and eluted with 65% EtOAc/Hex to afford the title compound (90 mg, 41%).


MS (ES) m/e 480.4 (M+1).

Example 22.

Preparation of 2-{1-Ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophene-6-carboxylic acid




A. {4-[1-Ethyl-1-(6-methoxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenoxy}-acetic acid methyl ester.




To a mixture of 4-[1-ethyl-1-(6-methoxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenol (2.53 g, 7.43 mmol) and acetonitrile (25 mL) is added methyl bromoacetate (0.84 mL, 8.92 mmol), powdered potassium carbonate (4.10 g, 29.72 mmol), and potassium iodide (0.12 g, 0.74 mmol). The resulting slurry is stirred at reflux temperature for 2 h, filtered, and concentrated. The residue is dissolved in diethyl ether (100 mL), washed with water (75 mL), brine (75 mL), dried over $MgSO_4$, filtered, and concentrated to give the title compound (1.5 g, 3.63 mmol, 49%). 1H NMR ($CDCl_3$), δ 0.74 (t, J = 7.2 Hz, 6H), 2.17 (q, J = 14.7, 7.5 Hz, 4H), 2.25 (s, 3H), 2.30 (s, 2H), 3.80 (s, 3H), 3.86 (s, 3H), 3.60 (d, J = 8.6 Hz, 1H), 6.89 (dd, J = 8.6, 2.5 Hz, 1H), 7.04 (s, 1H), 7.07 (s, 1H), 7.16 (d, J = 2.5 Hz, 1H), 7.55 (d, J = 8.9 Hz, 1H).

B. 3-{4-[1-Ethyl-1-(6-methoxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenoxy-methyl}-pentan-3-ol.



To a mixture of {4-[1-ethyl-1-(6-methoxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenoxy}-acetic acid methyl ester (1.50 g, 3.64 mmol) and THF (20 mL) is added ethyl magnesium bromide (3.0 mL, 9.1 mmol, 3M in THF). The solution is heated at a reflux for 3 h, diluted with saturated ammonia chloride solution (80 mL), extracted with diethyl ether (2 x 75 mL), dried over MgSO_4 , filtered, and concentrated. The residue is purified by silica gel chromatography (10% to 30% EtOAc gradient) to give the title compound (0.90g, 2.04 mmol, 56%). ^1H NMR (CDCl_3), δ 0.75 (t, $J = 7.6$ Hz, 6H), 0.95 (t, $J = 7.4$ Hz, 6H), 1.68 (q, $J = 14.8, 7.5$ Hz, 4H), 2.13-2.24 (m, 7H), 3.81 (s, 2H), 3.86 (s, 3H), 6.73 (dd, $J = 8.2$ Hz, 1H), 6.89 (dd, $J = 8.8, 2.6$ Hz, 1H), 7.06 (s, 2H), 7.09 (dd, $J = 8.4, 2.5$ Hz, 1H), 7.17 (d, $J = 2.7$ Hz, 1H), 7.55 (d, $J = 8.8$ Hz, 1H). LC/MS (m/z): calcd. for $\text{C}_{27}\text{H}_{37}\text{O}_3\text{S}$ ($\text{M}+\text{H}$) $^+$: 441.3; found: 441.2.


C. 2-{1-Ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl} benzo[b]thiophen-6-ol.



To a solution of 3-{4-[1-ethyl-1-(6-methoxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenoxy-methyl}-pentan-3-ol (0.80 g, 1.82 mmol) and DMF (10 mL) is added NaSEt (1.53 g, 18.15 mmol). The resulting solution is heated at 100°C overnight, diluted with diethyl ether (50 mL), washed with water (3 x 50 mL), brine


(50 mL), dried with MgSO_4 , filtered and concentrated. The resulting residue is purified by silica gel chromatography (10% to 40% EtOAc gradient) to give the title compound (0.67 g, 1.57 mmol, 87%). ^1H NMR (CDCl_3), δ 0.75 (t, $J = 7.2$ Hz, 6H), 0.95 (t, $J = 7.4$ Hz, 6H), 1.67 (q, $J = 15.1, 7.4$ Hz, 4H), 2.12-2.24 (m, 7H), 3.82 (s, 2H), 6.73 (d, $J = 8.6$ Hz, 1H), 6.79 (dd, $J = 9.0, 2.6$ Hz, 1H), 6.79 (dd, $J = 9.0, 2.6$ Hz, 1H), 7.00 (s, 1H), 7.05 (d, $J = 2.2$ Hz, 1H), 7.08 (dd, $J = 8.2, 2.6$ Hz, 1H), 7.11 (d, $J = 2.2$ Hz, 1H), 7.49 (d, $J = 9.0$ Hz, 1H). LC/MS (m/z): calcd. for $\text{C}_{26}\text{H}_{35}\text{O}_3\text{S}$ ($\text{M}+\text{H}$) $^+$: 427.2; found: 427.1.

D. Trifluoro-methanesulfonic acid 2-{1-ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophen-6-yl ester



To a solution of 2-{1-Ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl} benzo[b] thiophen-6-ol (0.67 g, 1.57 mmol) and dichloromethane (5 mL) is added NEt_3 (0.44 mL, 3.14 mmol). The solution is cooled to 0°C , and trifluoromethane-sulfonic anhydride (0.28 mL, 1.65 mmol) is added. The solution is warmed to RT, diluted with diethyl ether (30 mL), washed with saturated NaHCO_3 (20 mL), dried with MgSO_4 , filtered, and concentrated to give the title compound (0.22 g, 0.40 mmol, 85%). ^1H NMR (CDCl_3), δ 0.75 (t, $J = 7.3$ Hz, 6H), 0.95 (t, $J = 7.5$ Hz, 6H), 1.68 (q, $J = 14.8, 7.6$ Hz, 4H), 2.14-2.27 (m, 7H), 3.82 (s, 2H), 6.74 (d, $J = 8.3$ Hz, 1H), 7.03 (d, $J = 2.1$ Hz, 1H), 7.08 (dd, $J = 8.7, 2.5$ Hz, 1H), 7.13 (dd, $J = 8.7, 2.9$ Hz, 1H), 7.14 (s, 1H), 7.58 (d, $J = 2.1$ Hz, 1H), 7.70 (d, $J = 8.3$ Hz, 1H). LC/MS (m/z): calcd. for $\text{C}_{27}\text{H}_{34}\text{F}_3\text{O}_5\text{S}_2$ ($\text{M}+\text{H}$) $^+$: 559.2; found: 576.1.

E. 2-{1-Ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl} benzo[b]thiophene-6-carboxylic acid methyl ester.



To a solution of trifluoro-methanesulfonic acid 2-{1-ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl}-benzo[b]thiophen-6-yl ester (0.66 g, 1.19 mmol) and DMSO (15 mL) is added MeOH (10 mL), NEt₃ (0.77 mL), DPPB (0.054 g, 0.13 mmol), Pd(OAc)₂ (0.024, 0.01 mmol), and CO (100 psi). The solution is heated at 80 °C overnight, diluted with 1M HCl (100 mL), and extracted with diethyl ether (2 x 50 mL). The combined organic layers are washed with water (75 mL) then brine (75 mL), dried MgSO₄, filtered, and concentrated. The residue is purified by silica gel chromatography (10% to 40% EtOAc gradient) to give the title compound (0.84 g, 2.04 mmol, 68%). ¹H NMR (CDCl₃), δ 0.75 (t, *J* = 7.2 Hz, 6H), 0.95 (t, *J* = 7.7 Hz, 6H), 1.67 (q, *J* = 14.6, 7.5 Hz, 4H), 2.14-2.25 (m, 7H), 3.81 (s, 2H), 3.95 (s, 3H), 6.73 (d, *J* = 8.7 Hz, 1H), 7.04 (d, *J* = 2.2 Hz, 1H), 7.08 (dd, *J* = 8.4, 2.2 Hz, 1H), 7.20 (s, 1H), 7.71 (d, *J* = 8.5 Hz, 1H), 7.89 (dd, *J* = 8.4, 1.7 Hz, 1H), 8.39 (d, *J* = 1.7 Hz, 1H). LC/MS (*m/z*): calcd. for C₂₈H₃₇O₄S (M+H)⁺: 469.2; found: 486.2.


F. 2-{1-Ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl} benzo[b]thiophene-6-carboxylic acid.

2-{1-Ethyl-1-[4-(2-ethyl-2-hydroxy-butoxy)-3-methyl-phenyl]-propyl} benzo[b]thiophene-6-carboxylic acid methyl ester (0.48 g, 1.02 mmol) is dissolved in MeOH (2 mL) and treated with H₂O (0.5 mL) and NaOH (0.20 g, 5.12 mmol). The resulting mixture is heated at a reflux for 2 h, cooled to RT, and stirred overnight. The solution is diluted with H₂O (10 mL), adjusted to pH 3-4 using 1 M HCl, and extracted with EtOAc (40 mL). The EtOAc layer is washed with brine (20 mL), dried with MgSO₄, filtered, and concentrated to yield the title compound (0.42 g, 0.92 mmol, 89%). ¹H NMR (CDCl₃), δ 0.76 (t, *J* = 7.5 Hz, 6H), 0.95 (t, *J* = 7.5 Hz, 6H), 1.68 (q, *J* = 14.8, 7.5 Hz, 4H), 2.25-2.27 (m, 7H), 3.82 (s, 2H), 6.74 (d, *J* = 8.8 Hz, 1H), 7.05 (d, *J* = 2.3 Hz, 1H), 7.09 (dd, *J* = 8.5,

2.1 Hz, 1H), 7.24 (s, 1H), 7.75 (d, $J = 8.4$ Hz, 1H), 7.96 (dd, $J = 8.3, 1.7$ Hz, 1H), 8.48 (d, $J = 1.7$ Hz, 1H). Exact mass (m/z): calcd. for $C_{27}H_{35}O_4S$ ($M+H$)⁺: 454.2; found: 472.3.

Example 23.


Preparation of 5-{1-[4-(2-tert-Butyl-[1,3]dioxolan-2-ylmethoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid ethyl ester.



A solution of 5-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophene-2-carboxylic acid ethyl ester (170 mg, 0.354 mmol) in ethylene glycol (3.0 mL) is treated with $BF_3 \cdot Et_2O$ (2.0 mL). The mixture is stirred at 90 °C for 6 h, cooled to RT, and quenched with water (50 mL). The mixture is extracted with EtOAc (2 x 20 mL), and the organic layer is concentrated and purified using silica gel column chromatography (10% EtOAc/Hex) to afford the product as an oil (94 mg, 51%).
MS (ES) m/e 542.3 ($M+18$).

Example 24.

Preparation of methanesulfonic acid 2-{1-[4-(3,3-dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-1-ethyl-propyl}-benzo[b]thiophen-5-yl ester.



A solution of 1-{4-[1-ethyl-1-(5-hydroxy-benzo[b]thiophen-2-yl)-propyl]-2-methyl-phenoxy}-3,3-dimethyl-butan-2-one (260 mg, 0.612 mmol) in CH_2Cl_2 (20

mL) at RT is treated with Et₃N (0.40 mL) and methanesulfonyl chloride (140 mg, 1.22 mmol). The mixture is stirred for 30 min and quenched with water (1.0 mL). The organic layer is concentrated and purified using silica gel column chromatography (25% EtOAc/Hex) to afford the title compound (190 mg, 62%). NMR (ppm, CDCl₃), δ : 7.67 (1 H, d, $J=8.8$ Hz), 7.59 (1 H, d, $J=2.2$ Hz), 7.13 (1 H, dd, $J=2.2, 8.8$ Hz), 7.11 (1 H, s), 7.01-7.04 (2 H, m), 6.51 (1 H, d, $J=8.4$ Hz), 4.84 (2 H, s), 3.16 (3 H, s), 2.26 (3 H, s), 2.15 (4 H, q, $J=7.5$ Hz), 1.26 (9 H, s), 0.74 (6 H, t, $J=7.5$ Hz). MS (ES) m/e 520.3 (M+18).

Compounds of the Invention – Salts, Stereoisomers, & Prodrugs:

Salts of the compounds represented by formulae IA and IB are an additional aspect of the invention. The skilled artisan will also appreciate that the family of compounds of formulae I include acidic and basic members and that the present invention includes pharmaceutically acceptable salts thereof.

In those instances where the compounds of the invention possess acidic or basic functional groups various salts may be formed which are more water soluble and physiologically suitable than the parent compound. Representative pharmaceutically acceptable salts, include but are not limited to, the alkali and alkaline earth salts such as lithium, sodium, potassium, ammonium, calcium, magnesium, aluminum, zinc, and the like. Sodium and potassium salts are particularly preferred. Salts are conveniently prepared from the free acid by treating the acid in solution with a base or by exposing the acid to an ion exchange resin. For example, a carboxylic acid substituent on the compound of Formula I may be selected as -CO₂H and salts may be formed by reaction with appropriate bases (e.g., NaOH, KOH) to yield the corresponding sodium and potassium salt.

Included within the definition of pharmaceutically acceptable salts are the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention, for example, ammonium, quaternary ammonium, and amine cations, derived from nitrogenous bases of sufficient basicity to form salts with the compounds of this invention (see, for example, S. M. Berge, *et al.*, "Pharmaceutical Salts," *J. Phar. Sci.*, 66: 1-19 (1977)). Moreover, the basic group(s) of the compound

of the invention may be reacted with suitable organic or inorganic acids to form salts such as acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, camsylate, carbonate, chloride, choline, clavulanate, citrate, chloride, chloroprocaine, choline, diethanolamine, dihydrochloride, diphosphate, edetate, edisylate, estolate, esylate, ethylenediamine, fluoride, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, bromide, chloride, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, malseate, mandelate, meglumine, mesylate, mesviate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, palmitate, pamoate, pantothenate, phosphate, polygalacturonate, procaine, salicylate, stearate, subacetate, succinate, sulfate, tannate, tartrate, teoclate, tosylate, trifluoroacetate, trifluoromethane sulfonate, and valerate.

Certain compounds of the invention may possess one or more chiral centers and may thus exist in optically active forms. Likewise, when the compounds contain an alkenyl or alkenylene group there exists the possibility of cis- and trans- isomeric forms of the compounds. The R- and S- isomers and mixtures thereof, including racemic mixtures as well as mixtures of cis- and trans- isomers, are contemplated by this invention. Additional asymmetric carbon atoms can be present in a substituent group such as an alkyl group. All such isomers as well as the mixtures thereof are intended to be included in the invention. If a particular stereoisomer is desired, it can be prepared by methods well known in the art by using stereospecific reactions with starting materials which contain the asymmetric centers and are already resolved or, alternatively by methods which lead to mixtures of the stereoisomers and subsequent resolution by known methods. For example, a chiral column may be used such as those sold by Daicel Chemical Industries identified by the trademarks:

CHIRALPAK AD, CHIRALPAK AS, CHIRALPAK OD, CHIRALPAK OJ,
CHIRALPAK OA, CHIRALPAK OB, CHIRALPAK OC, CHIRALPAK OF,
CHIRALPAK OG, CHIRALPAK OK, and
CHIRALPAK CA-1.


By another conventional method, a racemic mixture may be reacted with a single enantiomer of some other compound. This changes the racemic form into a mixture of diastereomers. These diastereomers, because they have different melting points, different boiling points, and different solubilities can be separated by conventional means, such as crystallization.

The present invention is also embodied in mixtures of compounds of formulae IA or IB.


Prodrugs are derivatives of the compounds of the invention which have chemically or metabolically cleavable groups and become by solvolysis or under physiological conditions the compounds of the invention which are pharmaceutically active in vivo. Derivatives of the compounds of this invention have activity in both their acid and base derivative forms, but the acid derivative form often offers advantages of solubility, tissue compatibility, or delayed release in a mammalian organism (see, Bundgard, H., Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985). Prodrugs include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acidic compound with a suitable alcohol, or amides prepared by reaction of the parent acid compound with a suitable amine. Simple aliphatic or aromatic esters derived from acidic groups pendent on the compounds of this invention are preferred prodrugs. In some cases it is desirable to prepare double ester type prodrugs such as (acyloxy) alkyl esters or ((alkoxycarbonyl)oxy)alkyl esters. Particularly preferred esters to use as prodrugs are; methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, morpholinoethyl, and N,N-diethylglycolamido.

N,N-diethylglycolamido ester prodrugs may be prepared by reaction of the sodium salt of a compound of Formula IA or IB (in a medium such as dimethylformamide) with 2-chloro-N,N-diethylacetamide (available from Aldrich Chemical Co., Milwaukee, Wisconsin USA; Item No.25,099-6).


Morpholinylethyl ester prodrugs may be prepared by reaction of the sodium salt of a compound of Formula IA or IB (in a medium such as dimethylformamide) 4-(2-chloroethyl)morpholine hydrochloride (available from Aldrich Chemical Co., Milwaukee, Wisconsin USA, Item No. C5,220-3). For example, prodrugs may be prepared by reaction of the sodium salt for a compound of Formula IA or IB with;



and sodium iodide to provide the ester prodrug pendent group



Also, lower alkyl (viz., C₁-C₈) ester prodrugs may be prepared by conventional means such as reacting the sodium or potassium salt (derived by forming the salt of any acidic compound of the invention; viz., reaction of a base such as KOH with an acidic group such as -CO₂H) of a compound of Formula IA or IB with an alkyl iodide such as methyl iodide, ethyl iodide, n-propyl iodide, isopropyl iodide. Typical ester prodrug substituents are



Pharmaceutical Formulations containing the Novel Compounds of the Invention:

Pharmaceutical formulations of the invention are prepared by combining (e.g., mixing) a therapeutically effective amount of the compound of the invention (compounds of Formula IA or IB) together with a pharmaceutically acceptable carrier or diluent. The present pharmaceutical formulations are prepared by known procedures using well-known and readily available ingredients.

In making the compositions of the present invention, the compounds of the invention will usually be admixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), or ointment, containing, for example, up to 10% by weight of the compound. The compounds of the present invention are preferably formulated prior to administration.

The compounds of the invention may also be delivered by suitable formulations contained in a transderm patch. Alternatively, the compounds of the invention may be delivered to a patient by sublingual administration.

For the pharmaceutical formulations any suitable carrier known in the art can be used. In such a formulation, the carrier may be a solid, liquid, or mixture of a solid and a liquid. Solid form formulations include powders, tablets and capsules. A solid carrier can be one or more substances which may also act as flavoring agents, lubricants, solubilisers, suspending agents, binders, tablet disintegrating agents and encapsulating material.

Tablets for oral administration may contain suitable excipients such as calcium carbonate, sodium carbonate, lactose, calcium phosphate, together with disintegrating agents, such as maize, starch, or alginic acid, and/or binding agents, for example, gelatin or acacia, and lubricating agents such as magnesium stearate, stearic acid, or talc.

In powders the carrier is a finely divided solid which is in admixture with the finely divided Active ingredient. In tablets a compound of the invention I is mixed with a carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from about 1 to about 99 weight percent of the compound of this invention. Suitable solid carriers are magnesium carbonate, magnesium stearate, talc, sugar lactose, pectin, dextrin, starch, gelatin, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, low melting waxes, and cocoa butter.

Sterile liquid form formulations include suspensions, emulsions, syrups and elixirs.

The Active Ingredient may be dissolved or suspended in a pharmaceutically acceptable carrier, such as sterile water, sterile organic solvent or a mixture of both. The compounds can often be dissolved in a suitable organic solvent, for instance aqueous propylene glycol. Other compositions can be made by dispersing the finely divided compounds of the invention in aqueous starch or sodium carboxymethyl cellulose solution or in a suitable oil.

Methods of Using the Compounds of the Invention:

Many disease states are benefited by treatment with the compounds of Formula IA or IB include, but are not limited to:

- disease states characterized by abnormal calcium regulation
- disease states characterized by abnormal cell proliferation
- disease states characterized by abnormal cell differentiation

disease states characterized by abnormal immune response
disease states characterized by abnormal dermatological conditions
disease states characterized by neurodegenerative condition
disease states characterized by inflammation
disease states characterized by vitamin D sensitivity
disease states characterized by hyperproliferative disorders.

Specific disease states benefited by treatment of the compounds of Formula I and II include, but are not limited to:

Acne
Actinic keratosis
Alopecia
Alzheimer's disease
Bone maintenance in zero gravity
Bone fracture healing
Breast cancer
Chemoprevention of Cancer
Crohn's disease
Colon cancer
Type I diabetes
Host-graft rejection
Hypercalcemia
Type II diabetes
Leukemia
Multiple sclerosis
Myelodysplastic syndrome
Insufficient sebum secretion
Osteomalacia
Osteoporosis
Insufficient dermal firmness
Insufficient dermal hydration
Psoriatic arthritis
Prostate cancer
Psoriasis

Renal osteodystrophy

Rheumatoid arthritis

Scleroderma

Skin cancer

Systemic lupus erythematosus

Skin cell damage from Mustard vesicants

Ulcerative colitis

Vitiligo

Wrinkles

Particularly preferred is the treatment of psoriasis and osteoporosis by administration to a mammal (including a human) of a therapeutically effective amount of compounds of Formulae IA or IB.. By "pharmaceutically effective amount" it is meant that quantity of pharmaceutical agent corresponding to formulae IA or IB which prevents, removes or reduces the deleterious effects of a disease state in mammals, including humans.

The specific dose of a compound administered according to this invention to obtain therapeutic or prophylactic effects will, of course, be determined by the particular circumstances surrounding the case, including, for example, the compound administered, the route of administration and the condition being treated. Typical daily doses will contain a pharmaceutically effective amount typically in the range of from about 0.0001 mg/kg/day to about 50 mg/kg/day of body weight of an active compound of this invention. Preferably the dose of compounds of the invention will be from 0.0001 to 5 mg/kg/day of body weight.

Preferably compounds of the invention or pharmaceutical formulations containing these compounds are in unit dosage form for administration to a mammal. The unit dosage form can be a capsule or tablet itself, or the appropriate number of any of these. The quantity of Active ingredient in a unit dose of composition may be varied or adjusted from about 0.0001 to about 1000 milligrams or more according to the particular treatment involved. It may be appreciated that it is necessary to make routine variations to the dosage depending on the age and condition of the patient. Dosage will also depend on the route of administration. The compounds of the invention may be administered by a variety of routes including oral, aerosol, rectal, transdermal, sublingual, subcutaneous, intravenous, intramuscular, and intranasal. Particularly preferred is the treatment

of psoriasis with an ointment type formulation containing the compounds of the invention. The ointment formulation may be applied as needed, typically from one to 6 times daily.

Treatment of psoriasis is preferably done with topical application by a formulation in the form of a cream, oil, emulsion, paste or ointment containing a therapeutically effective amount of a compound of the invention.. The formulation for topical treatment contains from 0.5 to 0.00005 weight percent, preferably from .05 to 0.0005 weight percent, and most preferably from 0.025 to 0.001 of a Active Ingredient.

For example, two semisolid topical preparations useful as vehicles for VDR modulators in treatment and prevention of psoriasis are as follows:

Polyethylene Glycol Ointment USP (p. 2495)

Prepare Polyethylene Glycol Ointment as follows:

Polyethylene Glycol 3350	400 g.
Polyethylene Glycol 400	<u>600 g.</u>
To make	1000 g.

Heat the two ingredients on a water bath to 65C. Allow to cool, and stir until congealed. If a firmer preparation is desired, replace up to 100 g of the polyethylene glycol 400 with an equal amount of polyethylene glycol 3350.

Hydrophilic Ointment USP (p. 1216)

Prepare Hydrophilic Ointment as follows:

Methylparaben	0.25 g.
Propylparaben	0.15 g.
Sodium Lauryl Sulfate	10 g.
Propylene Glycol	120 g.
Stearyl Alcohol	250 g.
White Petrolatum	250 g.
Purified Water	<u>370 g.</u>
To make about	1000 g.

The Stearyl Alcohol and White Petrolatum are melted on a steam bath, and warmed to about 75C. The other ingredients, previously dissolved in the water are added, warmed to 75C, and the mixture stirred until it congeals.

For each of the above formulations the Active Ingredient is added during the

heating step in an amount that is from 0.5 to 0.00005 weight percent, preferably from .05 to 0.0005 weight percent, and most preferably from 0.025 to 0.001 weight percent of the total ointment weight. (Source: - United States Pharmacopoeia 24, United States Pharmacopeial Convention, 1999)

Conventional therapy for osteoporosis includes; (i) estrogens, (ii) androgens, (iii) calcium supplements, (iv) vitamin D metabolites, (v) thiazide diuretics, (vi) calcitonin, (vii) bisphosphonates, (viii) SERMS, and (ix) fluorides (see, Harrison's Principles of Internal Medicine, 13th edition, 1994, published by McGraw Hill Publ., ISBN 0-07-032370-4, pgs.2172-77; the disclosure of which is incorporated herein by reference.). Any one or combination of these conventional therapies may be used in combination with the method of treatment using compounds of Formulae IA or IB as taught herein. For example, in a method of treating osteoporosis, the vitamin D receptor modulator compounds of the invention may be administered separately or simultaneously with a conventional therapy. Alternatively, the vitamin D receptor modulator compounds of the invention may be combined with conventional therapeutic agents in a formulation for treatment of osteoporosis such as set out below:

A formulation for treating osteoporosis comprising:

Ingredient (A1): a vitamin D receptor modulator represented by formula (IA) or (IB), or a pharmaceutically acceptable salt or prodrug derivative thereof;

Ingredient (B1):

one or more co-agents that are conventional for treatment osteoporosis selected from the group consisting of:

- a. estrogens,
- b. androgens,
- c. calcium supplements,
- d. vitamin D metabolites,
- e. thiazide diuretics,
- f. calcitonin,
- g. bisphosphonates,
- h. SERMS, and
- i. fluorides.

Ingredient (C1): optionally, a carrier or diluent.

Typically useful formulations are those wherein the weight ratio of (A1) to (B1) is from 10:1 to 1:1000 and preferably from 1:1 to 1:100.

Combination Therapy for Psoriasis:

Conventional therapy for psoriasis includes topical glucocorticoids, salicylic acid, crude coal tar, ultraviolet light, and methotrexate (see, Harrison's Principles of Internal Medicine, 13th edition, 1994, published by McGraw Hill Publ., ISBN 0-07-032370-4, pgs.2172-77). Any one or combination of these conventional therapies may be used in combination with the method of treatment using compounds of Formulae IA or IB as taught herein. For example, in a method of treating osteoporosis, the vitamin D receptor modulator compounds of the invention (e.g., as defined by formula I) may be topically administered separately or simultaneously with a conventional therapy. Alternatively, the vitamin D receptor modulator compounds of the invention may be combined with conventional therapeutic agents in a topically applied formulation for treatment of osteoporosis such as set out below:

A formulation for treating psoriasis comprising:

Ingredient (A2): a vitamin D receptor modulator represented by formula (IA) or (IB), or a pharmaceutically acceptable salt or prodrug derivative thereof;

Ingredient (B2):

one or more co-agents that are conventional for treatment of psoriasis selected from the group consisting of:

- a. topical glucocorticoids ,
- b. salicylic acid, or
- c. crude coal tar.

Ingredient (C2): optionally, a carrier or diluent.

Typically useful formulations are those wherein the weight ratio of (A2) to (B2) is from 1:10 to 1:100000 and preferably from 1:100 to 1:10000.

Experimental Results:

Table 1

Summary of Experimental Results

Test Cmpd. ¹	RXR-VDR heterodimer ² EC ₅₀ (nM)	VDR EC ₅₀ (nM) (Caco-2 cells) ³	OCN Promoter ⁴ EC ₅₀ (nM)	Mouse Hypercal ⁵ µg/Kg/d
Ex. 1	234/14	432	2/1	<300
Ex. 2	30	411	1	
Ex. 3	442/67	869	7/168	
Ex. 4	401/19	555	20/15	
Ex. 5	2/27	342	1	>300
Ex. 6	18/0.4	331	1/0.1	
Ex 7	159/9	791	1	>300
Ex. 8	135/1	409	0.8/0.3	
Ex. 9	959/27	1271	41/35	
Ex. 10	323	690	24/18	
Ex. 11	45	299	0.2	>300
Ex. 12	19/2	137	0.3	
Ex. 13	488/111	994	9/28	>300
Ex 14	376/26	941	6	
Ex 15	539/63		20/60	<1000
Ex 16	254/108	938	19/39	
Ex 17	272/51	631	4/3	>300
Ex 18	242/3	748	2.4/1.3	
Ex 19	114/21	815	4.7	>300
Ex 20	24	314	5	
Ex 21	134/24	708	29/48	>1000

Ex 22	458/139	1057	32/15	
Ex 23			161	
Ex 24	695		215	

Table 2

Summary of Experimental Results

Test Cmpd. ¹	Kera. Prolif. IC ₅₀ (nM)	IL-10. IC ₅₀ (nM)
Ex. 1	11/23	109
Ex. 2	1	24
Ex. 3	>1000	351
Ex. 4		
Ex. 5	4	6
Ex. 6	4	4
Ex 7	1000	
Ex. 8	1	15
Ex. 9	1000	
Ex. 10	573	
Ex. 11	2	10
Ex. 12	1	
Ex. 13		
Ex 14		
Ex 15		
Ex 16	384	
Ex 17	404	

Ex 18	1.8	89
Ex 19	16	149
Ex 20	98	386
Ex 21	162	
Ex 22	154	
Ex 24	232	

Table 3 of Experimental Results
(comparison compounds)

Test Cmpd. ¹	RXR-VDR (SaOS-2 cells) ² EC ₅₀ (nM)	VDR CTF (Caco-2 cells) ³ EC ₅₀ (nM)	OCN Promoter ⁴ EC ₅₀ (nM)	Mouse Hypercal ⁵ µg/Kg/d
AA	5.02	16	5	0.06
BB	10.32	169.81	8.24	20
CC	2427.7		2680.9	
DD	109.44		31.1	1000
EE	429.99	891.16	341.25	1000
FF	3/1	57	0.28	

Table 4 of Experimental Results
(comparison compounds)

Test Cmpd. ¹	Kera. Prolif. IC ₅₀ (nM)	IL-10 IC ₅₀ (nM)
----------------------------	--	--------------------------------

AA	120	1.2
BB	10	28
CC	-	-
DD	1060	
EE		
FF	103	0.5

Explanation of Table 5 and 6 column numerical superscripts:

1. Test Compound numbers refer to the products of the corresponding

Example Nos. that is, compounds within the scope of the invention


2. The control experiments are done with the double letter coded compounds identified as follows:

“AA” = 1 α ,25-dihydroxyvitamin D₃


“BB” = 3-(4-{1-Ethyl-1-[4-(2-hydroxy-3,3-dimethyl-butoxy)-3-methyl-phenyl]-propyl}-2-methyl-phenoxy)-propane-1,2-diol

“CC” = 1-(4-{1-[4-(3,3-Dimethyl-2-oxo-butoxy)-3-methyl-phenyl]-cyclohexyl}-2-methyl-phenoxy)-3,3-dimethyl-butan-2-one


“DD” = compound represented by the formula:



“EE” = compound represented by the formula:



“FF” = calcipotriol (structural formula below):



2. The RXR-VDR heterodimerization (SaOS-2 cells) test is described in the “Assay” section of the Description, *infra*.
3. The VDR CTF (Caco-2 cells) test is described in the “Assay” section of the Description, *infra*.
4. The OCN Promoter test is described in the “Assay” section of the Description, *infra*.
5. The Mouse Hypercalcemia test is described in the “Assay” section of the Description, *infra*.
6. The keratinocyte proliferation assay is described in the “Assay” section of the Description, *infra*.
7. The IL-10 induction assay is described in the “Assay” section of the Description, *infra*.

Assay Methods

Use of the Assay Methods:

The evaluation of the novel compounds of the invention for osteoporosis and other related diseases is done using a plurality of test results. The use of multiple assays is necessary since the combined properties of (i) high activity for the vitamin D receptor, and (ii) prevention of hypercalcemia must be achieved to have utility for the methods of treating diseases, which are also, aspects of this invention. Some of the tests described below are believed related to other tests and measure related properties of compounds. Consequently, a compound may be considered to have utility in the

practice of the invention if it meets most, if not all, of the acceptance criteria for the above described tests.

The evaluation of the novel compounds of the invention for psoriasis is done using the Keratinocyte Proliferation Assay in combination with other assays that measure inhibition of IL-2 production and stimulation of IL-10 production in peripheral blood mononuclear cells (PBMCs).

Brief Description, Utility and Acceptance Criteria for the Assay Methods:

1. The RXR-VDR heterodimer Assay:

This assay provides the VDR activity of a test compound. It is desirable to have low EC₅₀ values for a compound in this assay. The lower the EC₅₀ value, the more active the compound will be as a VDR agonist. Desired assay results are EC₅₀ values less than or equal to 600 nM. Preferred assay results are less than 250 nM, and most preferably less than 150 nM.

2. The Caco-2 cell Co-transfection Assay:

The Caco-2 cell assay is an indicator for the undesirable condition of hypercalcemia. This co-transfection assay is a surrogate assay for in vivo calcemic activity of VDR ligands. It is desirable to have high EC₅₀ values for a test compound in this assay. The higher the EC₅₀ values for a compound the less calcemic it will be in vivo. Desired assay results are EC₅₀ greater than or equal to 300 nM. Preferred assay results are greater than 1000 nM.

3. The OCN (osteocalcin) Promoter Assay

The OCN Promoter Assay is an indicator and marker for osteoporosis. Desired assay results are EC₅₀ less than or equal to 325 nM. Preferred assay results are less than 50 nM.

4. The Mouse Hypercalcemia Assay

The Mouse Hypercalcemia Assay is a six day hypercalcemia test for toxicity and selectivity. Acceptable test results are levels greater than 300 µg/kg/day. Preferred assay results are levels greater than 1000 µg/kg/day.

5. The Keratinocyte Proliferation Assay

This Assay is indicative for the treatment of psoriasis. An acceptable test result is IC50 value of less than or equal to 300 nM. Preferred assay results are IC50 values of less than 100 nM.

6. The IL-10 induction Assay

This is an in vitro efficacy assay for psoriasis, abscess and adhesion. Psoriasis involves both keratinocytes and immune cells. IL-10 is a unique cytokine because it is anti-inflammatory and immunosuppressive. This assay tells us whether a VDRM is able to function as an agonist in PBMCs (primary blood mononuclear cells) or not. A lower EC50 value is desirable in this assay since a compound with a lower EC50 value will be a better agonist in PBMCs. An acceptable test result is an EC50 value of less than 200 nM. Preferred assay results are EC50 values of less than 100 nM.

7. Other Compound Assay Standards

An alternative measure of the therapeutic index (bone efficacy vx. Hypervcalcemia) of compounds of the invention for treatment of osteoporosis is a numerical ratio calculated as follows:

Dose Threshold needed to induce hypercalcemia

divided by

Dose Threshold needed for bone efficacy

An alternative measurse of the therapeutic index (in vivo keratinocyte proliferation vs. hypercalcemia) of compounds of the invention for treatment of psoriasis is a numerical ratio calculated as follows:

Dose Threshold needed to induce hypercalcemia

divided by

Dose Threshold needed to induce keratinocyte proliferation

For the above ratios, Dose Thresholds are determined from dose response curve data.

Details of the Assay Methods:(1) Materials and Method for RXR-VDR Heterodimerization Assay:

Transfection Method:

- FuGENE 6 Transfection Reagent (Roche Cat # 1 814 443)

Growth Media:

- D-MEM High Glucose (Gibco BRL Cat # 11054-020), 10% FBS, 1% antibiotic-antimycotic (Ab-Am)

FBS heat inactivated (Gibco BRL Cat # 10092-147)

Ab-Am (Gibco BRL Cat # 15240-062)

Cells:

- Grow SaOs-2 cells in T-152 cm² culture flasks in *growth media*.
- Keep the density at 5-6 x 10⁵ cells/ml
- Passage cells 1:3 twice a week
- Add Trypsin EDTA (Gibco BRL Cat # 25300-020) and incubate
- Resuspend cells in plating media and transfer into growth media.

Wash Media:

- HBSS Low Glucose Without Phenol Red (Gibco BRL Cat # 14175-095), 1% Ab-Am

Plating Media:

- D-MEM Low Glucose Without Phenol Red (Gibco BRL Cat # 11054-020), 1% Ab-Am

D-MEM

Stripped FBS (Hyclone Cat# SH30068.03 Lot # AHM9371)

Ab-Am

Transfection / Treatment Media:

- D-MEM Low Glucose Without Phenol Red only

T-152 cm² culture flask:

- Use Corning Costar T-152 cm² culture flask (Cat # 430825) to grow the cells

Flat well Plates:

- Use well plate to plate cells
- Use Deep well plate sterile to make up treatment media.

Luciferase Assay Reagent:

- Use Steady-Glo Luciferase Reagent from Promega (Cat # E2550) Consists of:

- a. E2533 Assay Substrate, lyophilized product and

- b. E2543 Assay Buffer.

- Thaw at room temperature

- Store

Cell Harvesting

Aspirate media from culture flask, rinse cells with HBSS and aspirate.

Add trypsin and incubate.

When cells appear detached, resuspend cells in *growth media*.

Transfer into a new flask with fresh *growth media* for passaging the cells.

Plate well plates and two extra plates

A. Cell Count

Mix the cell suspension using pipette

Use *Hematocytometer* to count the cells

Load cell suspension onto the hemocytometer chamber

Count cells.

Plate seeding:

Use plating media 10 % Stripped FBS in D-MEM Low Glucose, Without Phenol Red,

1% Ab-Am

Plate 14 plates @ 165 μ l / well.

In sterile flask add cell suspension

to plating media.

Mix .

Add cells / well.

Place the cells in the incubator.

Cells should be about 75 % confluent prior to transfection.

DAY 2: Transfection

Step 1: DNA and Media

Add plain DMEM media to tubes for mixing the DNA

Add the Reporter gene pFR-LUC

Add the Gal4-RXR-DEF and VP16-VDR-LBD

Step 2: FuGENE and Media

Prepare plain DMEM media in a tubes for mixing FuGENE

Add *FuGENE 6 Transfection Reagent*

Incubate

Step 3: FuGENE , DNA and Media Complex

Add FuGENE Media complex from step 2 to DNA Media complex from step 1

Incubate

Step 4: FuGENE , DNA and Media Complex to-well plate

Add FuGENE-DNA-Media complex from step 3 to each plate

Incubate.

Day 3: Dosing

Treatment preparation

Allow for transfection time

Make a stock solution of the compounds in DMSO

Vortex until all the compounds has been dissolved.

Further dilute in D-MEM (Low Glucose – With out Phenol Red)

Add compounds in quadruplicate to give final volume

Incubate.

Day 4: Luciferase Assay

Read the plates after drug treatment

Remove part of media from all the wells and leave remainder

Add Steady-Glo Luciferase Reagent mixture / wells

Incubate

Count each well using a Luminescence counter, Top Count NXT by Packard

Set a delay between plates to reduce the background.

(2) Materials and Method for The Caco-2 Cell Assay:

Caco-2 cells, grown in phenol red free, DMEM (Invitrogen, Carlsbad, CA) containing 10 % charcoal-stripped FCS (Hyclone, Logan, UT), were transfected with Fugene 6 reagent (Roche Diagnostics, Indianapolis, IN).

Cells (5000/well) were plated 18 h before transfection in a 96 well plate. The

Cells were transfected with Gal4-responsive reporter pFRLuc (150 ng, Stratagene, La Jolla CA) and the receptor expression vector pGal4-VDR-LBD (10 ng), along with Fugene 6 reagent (0.2 µl/well). The DNA-Fugene complex was formed by incubating the mixture for 30 min at room temperature. The cells were transfected in triplicate for 5 h, and treated with various concentrations of VDR ligands (from 0.01 nM to 10,000 nM concentration range) 18h post-transfection. The luciferase activity was quantified using Steady-Glo reagent kit (Promega, Madison, WI) as per manufacturer's specifications.

(3) Materials and Method for The OCN Promoter Assay:

The activation of osteocalcin by VDR ligands was evaluated in a rat osteoblast-like cell line RG-15 (ROS 17/2.8) stably expressing rat osteocalcin promoter fused with luciferase reporter gene. The stable cell lines were established as reported before (Activation of Osteocalcin Transcription involves interaction of protein kinase A- and Protein kinase C-dependent pathways. Boguslawski, G., Hale, L. V., Yu, X.-P., Miles, R. R., Onyia, J. E., Santerre R. F., Chandrasekhar, S. J Biol. Chem. 275, 999-1006, 2000). Confluent RG-15 cells maintained in DMEM/F-12 medium (3:1) containing 5% FBS, 300 µg/ml G418 and at 37°C under 5% CO₂/95% air atmosphere were trypsinized (0.25% trypsin) and plated into white opaque 96-well cell culture plates (25000 cells/well). After 24 hr, cells (in DMEM/F-12 medium + 2% FBS) were treated with various concentrations of compounds, dissolved in DMSO. The final DMSO concentration remained at 0.01% (v/v). After 48 hr treatment, the medium was removed, cells were lysed with 50 µl of lysis buffer (From Luciferase reporter assay system, Roche Diagnostics, Indianapolis, IN) and assayed for luciferase activity using the Luciferase Reporter Gene Assay kit from Boehringer Mannheim as per manufacturer's specifications.

(4) Materials and Method for The Mouse Hypercalcemia Assay:

Weanling, virus -antibody-free, five to six weeks old female DBF mice (Harlan, Indianapolis, IN) are used for all the studies. Animals are allowed to

acclimate to local vivarium conditions for 2 days. Mice are maintained on a 12 hr light/dark cycle at 22°C with ad lib access to food (TD 5001 with 1.2% Ca and 0.9%P, Teklad, Madison, WI) and water. The animals then are divided into groups with 4-5 mice per group. Different doses of test compounds prepared in 10% Ethanol and 90% sesame oil are administered to mice orally via gavage for 6 days. 1α - $25(\text{OH})_2\text{D}_3$ 0.5µg/kg/d was also given to one group of mice as the positive control. Serum ionized calcium is evaluated at 6 hours after the last dosing under isoflurane anesthesia by Ciba-Corning Ca^{++}/PH Analyzer, (Model 634, Chiron Diagnostics Corp., East Walpole, MA). Raw data of group differences is assessed by analysis of variance (ANOVA) using Fisher's protected least significant difference (PLSD) where the significance level was $P < 0.05$.

(5) The Keratinocyte Proliferation Assay:

KERtr cells (Human skin keratinocyte transformed with a retrovirus vector, obtained from ATCC) were plated in 96-well flat-bottomed plates (3000 cells/well) in 100 µl keratinocyte serum free medium supplemented with bovine pituitary extract in the absence of EGF (Life Technologies, Rockville, MD) and incubated at 37°C for two days. The cells were treated with various concentrations of VDR ligands (ten-fold serial dilution from 10,000 nM to 0.1 nM in triplicate), dissolved in 100 µl keratinocyte serum free medium supplemented with bovine pituitary extract in the absence of EGF and incubated at 37°C for 72hr. BrdU (5-bromo-2'-deoxyuridine) incorporation was analyzed as a measure of DNA replication (Cell proliferation ELISA kit, Roche Diagnostics, Indianapolis, IN) and absorbance was measured at 405 nm. Potency values (IC_{50}) values were determined as the concentration (nM) of compound that elicited a half-maximal response.

(6) Materials and Method for human IL-10 Induction Assay:

Isolation of peripheral blood mononuclear cells (PBMCs):

- A. Collect 50 ml of human blood and dilute with media, RPMI-1640.
- B. Prepare sterile tubes with ficol.
- C. Add diluted blood to tubes.
- D. Centrifuge.
- E. Discard the top layer and collect the cells from middle layer.

F. Divide all cells into four tubes and add media.

G. Centrifuge.

H. Aspirate off media and resuspend.

I. Collect all cells

J. Centrifuge. at 1200 rpm for 10 minutes.

K. Resuspend in RPMI-1640 with 2% FBS and count cells

Stimulation of PBMC:

L. Prepare TPA in DMSO.

M. Dissolve PHA in water .

N. Plate TPA/PHA treated PBMCs in well plates.

O. Incubate.

Treatment:

P. Prepare all compound dilutions in plain RPMI- 1640 media.

Q. Add diluted compound.

R. Incubate.


Sample Collection and assay:

S. Remove all the cells by centrifugation and assay the supernatant for IL-10 by immunoassay.

1) T. Perform IL-10 assay using anti-human IL-10 antibody coated beads, as described by the manufacturer (Linco Research Inc., St. Charles, MO).

WE CLAIM:

1. A compound or a pharmaceutically acceptable salt or a prodrug derivative thereof represented by formula (IA):




wherein



R and R' are independently C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, or together R and R' form a substituted or unsubstituted, saturated or unsaturated carbocyclic ring having from 3 to 8 carbon atoms;

RP₃ and RB are independently selected from hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, -O-C₁-C₅ alkyl, -S-C₁-C₅ alkyl, -O-C₁-C₅ fluoroalkyl, -CN, -NO₂, acetyl, -S-C₁-C₅ fluoroalkyl, C₂-C₅ alkenyl, C₃-C₅ cycloalkyl, or C₃-C₅ cycloalkenyl;

RP, RT₃, and RB' are independently selected from hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, -O-C₁-C₅ alkyl, -S-C₁-C₅ alkyl, -O-C₁-C₅ fluoroalkyl, -CN, -NO₂, acetyl, -S-C₁-C₅ fluoroalkyl, C₂-C₅ alkenyl, C₃-C₅ cycloalkyl, or C₃-C₅ cycloalkenyl;


(L_{P1}), (L_{P2}), and (L_{TB}) are divalent linking groups independently selected from the group consisting of








R40

R40



R40



where m is 0, 1, or 2, and each R40 is independently hydrogen, C₁-C₅ alkyl, or C₁-C₅ fluoroalkyl;

Z_p is

branched C₃-C₅ alkyl,

3-methyl-3-hydroxypentyl,

3-methyl-3-hydroxypentenyl,

3-methyl-3-hydroxypentynyl,

3-ethyl-3-hydroxypentyl,

3-ethyl-3-hydroxypentenyl,

3-ethyl-3-hydroxypentynyl,

3-ethyl-3-hydroxy-4-methylpentyl,

3-ethyl-3-hydroxy-4-methylpentenyl,

3-ethyl-3-hydroxy-4-methylpentynyl,

3-propyl-3-hydroxypentyl,

3-propyl-3-hydroxypentenyl,

3-propyl-3-hydroxypentynyl,

1-hydroxy-2-methyl-1-(methylethyl)propyl,

3-methyl-3-hydroxy-4,4-dimethylpentyl,
3-methyl-3-hydroxy-4,4-dimethylpentenyl,
3-methyl-3-hydroxy-4,4-dimethylpentynyl,
3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
3-ethyl-3-hydroxy-4,4-dimethylpentenyl,
3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
1-hydroxycyclopentenyl,
1-hydroxycyclohexenyl,
1-hydroxycycloheptenyl,
1-hydroxycyclooctenyl,
1-hydroxycyclopropyl,
1-hydroxycyclobutyl,
1-hydroxycyclopentyl,
1-hydroxycyclohexyl,
1-hydroxycycloheptyl, or
1-hydroxycyclooctyl;

provided, however, that when

Z_p is

3-methyl-3-hydroxypentyl,
3-methyl-3-hydroxypentenyl,
3-methyl-3-hydroxypentynyl,
3-ethyl-3-hydroxypentyl,
3-ethyl-3-hydroxypentenyl,
3-ethyl-3-hydroxypentynyl,
3-ethyl-3-hydroxy-4-methylpentyl,
3-ethyl-3-hydroxy-4-methylpentenyl,
3-ethyl-3-hydroxy-4-methylpentynyl,
3-propyl-3-hydroxypentyl,
3-propyl-3-hydroxypentenyl,
3-propyl-3-hydroxypentynyl,
3-methyl-3-hydroxy-4,4-dimethylpentyl,
3-methyl-3-hydroxy-4,4-dimethylpentenyl,
3-methyl-3-hydroxy-4,4-dimethylpentyl,

3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentenyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentynyl, or
 1-hydroxy-2-methyl-1-(methylethyl)propyl;

then (L_{P1}) and (L_{P2}) combine as a bond;

Z_{TB} is selected from

-O-(C₁-C₅ alkyl),
 -O-(C₂-C₅ alkenyl),
 -O-(C₃-C₅ cycloalkyl),
 -O-(C₃-C₅ cycloalkenyl),
 -O-(C₁-C₅ hydroxyalkyl),
 -O-(C₁-C₅ fluoroalkyl),
 -O-(C₁-C₅ alkyl)-phenyl,
 -O-(C₁-C₅ alkyl)-(O)-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl) NH₂,
 -O-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl)₂,
 -O-(C₁-C₅ alkyl)-C(O)-NH₂,
 -O-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
 -O-(C₁-C₅ alkyl)-C(O)-OH,
 -O-(C₁-C₅ alkyl)-C(O)-NH-5-tetrazolyl,
 -O-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl)-NH₂,
 -O-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
 -O-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
 -O-(C₁-C₅ alkyl)-N-pyrrolidine,
 -O-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),

-O-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-SO₂-NH₂,
-O-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-NH₂,
-O-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-5-tetrazolyl,
-O-CH₂-CO₂H,
-O-CH₂-5-tetrazolyl,
-O-(C₁-C₅ alkyl),
-O-C(O)-NH₂,
-O-C(O)-N-(CH₃)₂,
-O-C(S)-N-(CH₃)₂,
-O-C(O)-O-(C₁-C₅ alkyl),
-O-(5-tetrazolyl),
-O-SO₂-(C₁-C₅ alkyl),
-O-SO₂-NH₂,
-O-SO₂-NH-(C₁-C₅ alkyl),
-O-SO₂-N-(C₁-C₅ alkyl)₂,
-O-S(O)-(C₁-C₅ alkyl),
-O-S(O)-NH₂,
-O-S(O)-NH-(C₁-C₅ alkyl),
-O-S(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl),
-S-(C₂-C₅ alkenyl),

-S-(C₃-C₅ cycloalkyl),
-S-(C₃-C₅ cycloalkenyl),
-S-(C₁-C₅ fluoroalkyl),
-S-(C₁-C₅ hydroxyalkyl),
-S-(C₁-C₅ alkyl)-phenyl,
-S-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-OH,
-S-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-NH₂,
-S-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl) NH₂,
-S-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-S-(C₁-C₅ alkyl)-N-pyrrolidine,
-S-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-S-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-SO₂-NH₂,
-S-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-5-tetrazolyl,
-S-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-S(O)-NH₂,
-S-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),

-S-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₂-C₅ alkenyl),
-SO₂-(C₃-C₅ cycloalkyl),
-SO₂-(C₃-C₅ cycloalkenyl),
-SO₂-(C₁-C₅ hydroxyalkyl),
-SO₂-(C₁-C₅ fluoroalkyl),
-SO₂-(C₁-C₅)-phenyl,
-SO₂-NH₂,
-SO₂-NH-(C₁-C₅ alkyl),
-SO₂-NH-CH₂-C(O)OH,
-SO₂-NH-CH₂-C(O)(O-C₁-C₅ alkyl),
-SO₂-NH-(C₁-C₅ alkyl)-C(O)OH,
-SO₂-NH-(C₁-C₅ alkyl)-C(O)(O-C₁-C₅ alkyl),
-SO₂-NHC(O)-(C₃-C₆ cycloalkyl),
-SO₂-NH-C(O)-(C₁-C₅ alkyl),
-SO₂-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl) NH₂,
-SO₂-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-C(O)-NH₂,
-SO₂-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-SO₂-(C₁-C₅ alkyl)-N-pyrrolidine,

$\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)-(1-methylpyrrolidin-2-one-3-yl)}$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)-C(O)-O-(C}_1\text{-C}_5\text{ alkyl)}$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)-C(O)-OH}$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)-5-tetrazolyl}$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)}$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)-SO}_2\text{-NH}_2$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)-SO}_2\text{-NH-(C}_1\text{-C}_5\text{ alkyl)}$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)-SO}_2\text{-N-(C}_1\text{-C}_5\text{ alkyl)}_2$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)}$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)-P(O)-(O-C}_1\text{-C}_5\text{ alkyl)}_2$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ alkyl)}$,
 $\text{-SO}_2\text{-(C}_2\text{-C}_5\text{ alkenyl)}$,
 $\text{-SO}_2\text{-(C}_3\text{-C}_5\text{ cycloalkyl)}$,
 $\text{-SO}_2\text{-(C}_3\text{-C}_5\text{ cycloalkenyl)}$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ hydroxyalkyl)}$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{ fluoroalkyl)}$,
 $\text{-SO}_2\text{-(C}_1\text{-C}_5\text{)-phenyl}$,
 $\text{-SO}_2\text{-N=CHN(C}_1\text{-C}_5\text{ alkyl)}_2$,
 -S(O)-NH_2 ,
 $\text{-S(O)-NH-(C}_1\text{-C}_5\text{ alkyl)}$,
 $\text{-S(O)-NH-CH}_2\text{-C(O)OH}$,
 $\text{-S(O)-NH-(C}_1\text{-C}_5\text{ alkyl)-C(O)OH}$,
 $\text{-S(O)-NH-CH}_2\text{-C(O)(O-C}_1\text{-C}_5\text{ alkyl)}$,
 $\text{-S(O)-NH-(C}_1\text{-C}_5\text{ alkyl)-C(O)(O-C}_1\text{-C}_5\text{ alkyl)}$,
 $\text{-S(O)HC(O)-(C}_3\text{-C}_6\text{ cycloalkyl)}$,
 $\text{-S(O)-NH-C(O)-(C}_1\text{-C}_5\text{ alkyl)}$,
 $\text{-S(O)-N-(C}_1\text{-C}_5\text{ alkyl)}_2$,
 $\text{-S(O)-(C}_1\text{-C}_5\text{ alkyl)-O-(C}_1\text{-C}_5\text{ alkyl)}$,
 $\text{-S(O)-(C}_1\text{-C}_5\text{ alkyl)-C(O)-(C}_1\text{-C}_5\text{ alkyl)}$,

-S(O)-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-C(O)-NH₂,
-S(O)-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-NH-S(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-S(O)-(C₁-C₅ alkyl)-N-pyrrolidine,
-S(O)-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-S(O)-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-OH,
-S(O)-(C₁-C₅ alkyl)-5-tetrazolyl,
-S(O)-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-SO₂-NH₂,
-S(O)-(C₁-C₅ alkyl)-S(O)-NH₂,
-S(O)-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-S(O)-N=CHN(C₁-C₅ alkyl)₂,
-NHC(S)NH₂,
-NHC(S)NH-(C₁-C₅ alkyl),
-NHC(S)N-(C₁-C₅ alkyl)₂,

-NHC(S)NH-(C₂-C₅ alkenyl),
-NHC(S)NH-(C₃-C₅ cycloalkyl),
-NHC(S)NH-(C₃-C₅ cycloalkenyl),
-NHC(S)NH-(C₁-C₅ fluoroalkyl),
-NHC(S)NH-C₁-C₅ hydroxyalkyl,
-NHC(S)NH-(C₁-C₅ fluoroalkyl)
-NHC(S)NH-phenyl,
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-OH,
-NHC(S)NH-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-NH-S(O)-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-NHC(S)NH-(C₁-C₅ alkyl)-N-pyrrolidine,
-NHC(S)NH-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-
3-yl),
-NHC(S)NH-(C₁-C₅ alkyl)-5-tetrazolyl,
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),

-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-NH₂,
 -NHC(S)NH-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
 -NHC(S)NH-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
 -NHC(O)NH₂,
 -NHC(O)NH-(C₁-C₅ alkyl),
 -NHC(O)N-(C₁-C₅ alkyl)₂,
 -NHC(O)NH-(C₂-C₅ alkenyl),
 -NHC(O)NH-(C₃-C₅ cycloalkyl),
 -NHC(O)NH-(C₃-C₅ cycloalkenyl),
 -NHC(O)NH-(C₁-C₅ hydroxyalkyl),
 -NHC(O)NH-(C₁-C₅ fluoroalkyl),
 -NHC(O)NH-phenyl,
 -NHC(O)NH-(C₁-C₅ alkyl)-NH₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-NH₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-C(O)-NH₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
 -NHC(O)NH-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
 -NHC(O)NH-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
 -NHC(O)NH-(C₁-C₅ alkyl)-N-pyrrolidine,
 -NHC(O)NH-(C₁-C₅ alkyl)-
 (1-methylpyrrolidin-2-one-3-yl),

-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-OH,
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-5-tetrazolyl,
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-NH₂,
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₁-C₅ alkyl)-P(O)-O-(C₁-C₅ alkyl)₂ ,
-NH₂,
-NH-(C₁-C₅ alkyl),
-NH-CH₂-C(O)OH,
-N-(C₁-C₅ alkyl)₂,
-NH-C(O)-NH₂,
-NH-C(O)-NH-(C₁-C₅ alkyl),
-NH-C(O)-N-(C₁-C₅ alkyl)₂,
-NH-C(O)-(C₁-C₅ alkyl),
-NH-SO₂-(C₁-C₅ alkyl),
-NH-S(O)-(C₁-C₅ alkyl),
-N(CH₃)(OCH₃),
-N(OH)(CH₃),
-N-pyrrolidin-2-one,
-N-pyrrolidine,
-(1-methylpyrrolidin-2-one-3-yl),
-CO₂H,
-CO₂Me,
-CO₂Et,
-C(O)CH₂S(O)Me,
-C(O)CH₂S(O)Et,
-C(O)CH₂S(O)₂Me,
-C(O)CH₂S(O)₂Et,

-C(O)CH₂CH₂S(O)Me,
-C(O)CH₂CH₂S(O)Et,
-C(O)CH₂CH₂S(O)₂Me,
-C(O)CH₂CH₂S(O)₂Et,
-C(O)CH(Me)CH₂CO₂H,
-C(O)CH(Me)CH₂CO₂Me,
-C(O)CH(Me)CH₂CO₂Et,
-C(O)CH(Me)CH₂CO₂iPr,
-C(O)CH(Me)CH₂CO₂tBu,
-C(O)CH(Me)CH(Me)CO₂H,
-C(O)CH(Me)CH(Me)CO₂Me,
-C(O)CH(Me)CH(Me)CO₂Et,
-C(O)CH(Me)CH(Me)CO₂iPr,
-C(O)CH(Me)CH(Me)CO₂tBu,
-C(O)CH(Me)C(Me)₂CO₂H,
-C(O)CH(Me)C(Me)₂CO₂Me,
-C(O)CH(Me)C(Me)₂CO₂Et,
-C(O)CH(Me)C(Me)₂CO₂iPr,
-C(O)CH(Me)C(Me)₂CO₂tBu,
-C(O)CH(Me)CH(Et)CO₂H,
-C(O)CH(Me)CH(Et)CO₂Me,
-C(O)CH(Me)CH(Et)CO₂Et,
-C(O)CH(Me)CH(Et)CO₂iPr,
-C(O)CH(Me)CH(Et)CO₂tBu,
-C(O)C(O)OH,
-C(O)C(O)NH₂,
-C(O)C(O)NHMe,
-C(O)C(O)NMe₂,
-C(O)NH₂,




-C(O)NMe₂,
-C(O)NH-CH₂-C(O)OH,
-C(O)NH-CH₂-C(O)OMe,
-C(O)NH-CH₂-C(O)OEt,
-C(O)NH-CH₂-C(O)OiPr,
-C(O)NH-CH₂-C(O)OtBu,
-C(O)NH-CH(Me)-C(O)OH,
-C(O)NH-CH(Me)-C(O)OMe,
-C(O)NH-CH(Me)-C(O)OEt,
-C(O)NH-CH(Me)-C(O)iPr,
-C(O)NH-CH(Me)-C(O)tBu,
-C(O)NH-CH(Et)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OMe,
-C(O)NH-C(Me)₂-C(O)OEt,
-C(O)NH-C(Me)₂-C(O)iPr,
-C(O)NH-C(Me)₂-C(O)tBu,
-C(O)NH-CMe(Et)-C(O)OH,
-C(O)NH-CH(F)-C(O)OH,
-C(O)NH-CH(CF₃)-C(O)OH,
-C(O)NH-CH(OH)-C(O)OH,
-C(O)NH-CH(cyclopropyl)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-CF(Me)-C(O)OH,
-C(O)NH-C(Me)(CF₃)-C(O)OH,
-C(O)NH-C(Me)(OH)-C(O)OH,
-C(O)NH-C(Me)(cyclopropyl)CO₂H
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH₂-C(O)OMe,
-C(O)NMe-CH₂-C(O)OEt,
-C(O)NMe-CH₂-C(O)OiPr,
-C(O)NMe-CH₂-C(O)tBu,











-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH(Me)-C(O)OH,
-C(O)NMe-CH(F)-C(O)OH,
-C(O)NMe-CH(CF₃)-C(O)OH,
-C(O)NMe-CH(OH)-C(O)OH,
-C(O)NMe-CH(cyclopropyl)-C(O)OH,
-C(O)NMe-C(Me)₂-C(O)OH,
-C(O)NMe-CF(Me)-C(O)OH,
-C(O)NMe-C(Me)(CF₃)-C(O)OH,
-C(O)NMe-C(Me)(OH)-C(O)OH,
-C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,
-C(O)NHS(O)Me,
-C(O)NHSO₂Me,
-C(O)-NH-5-tetrazolyl,
-C(O)NHS(O)Me,
-C(O)NHS(O)Et,
-C(O)NHSO₂Me,
-C(O)NHSO₂Et,
-C(O)NHS(O)iPr,
-C(O)NHSO₂iPr,
-C(O)NHS(O)tBu,
-C(O)NHSO₂tBu,
-C(O)NHCH₂S(O)Me,
-C(O)NHCH₂S(O)Et,
-C(O)NHCH₂SO₂Me,
-C(O)NHCH₂SO₂Et,
-C(O)NHCH₂CH₂S(O)Me,
-C(O)NHCH₂CH₂S(O)Et,
-C(O)NHCH₂CH₂SO₂Me,
-C(O)NHCH₂CH₂SO₂Et,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)SO₂Me,

-C(O)-N(Me)-5-tetrazolyl,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)S(O)Et,
-C(O)N(Me)SO₂Me,
-C(O)N(Me)SO₂Et,
-C(O)N(Me)S(O)iPr,
-C(O)N(Me))SO₂iPr,
-C(O)N(Me))S(O)tBu,
-C(O)N(Me)SO₂tBu,
-C(O)N(Me)CH₂S(O)Me,
-C(O)N(Me)CH₂S(O)Et,
-C(O)N(Me)CH₂SO₂Me,
-C(O)N(Me)CH₂SO₂Et,
-C(O)N(Me)CH₂CH₂S(O)Me,
-C(O)N(Me)CH₂CH₂S(O)Et,
-C(O)N(Me)CH₂CH₂SO₂Me,
-C(O)N(Me)CH₂CH₂SO₂Et,
-CH₂CO₂H,
-CH₂-5-tetrazolyl,
-CH₂CO₂Me,
-CH₂CO₂Et,
-CH₂NHS(O)Me,
-CH₂NHS(O)Et,
-CH₂NHSO₂Me,
-CH₂NHSO₂Et,
-CH₂NHS(O)iPr,
-CH₂NHSO₂iPr,
-CH₂NHS(O)tBu,
-CH₂NHSO₂tBu,
-CH₂NHCH₂CH₂SO₂CH₃,

-CH₂NH(CH₂CO₂H),
-CH₂N(C(O)Me)(CH₂CO₂H),
-CH₂-N-pyrrolidin-2-one,
-CH₂-(1-methylpyrrolidin-2-one-3-yl),
-CH₂S(O)Me,
-CH₂S(O)Et,
-CH₂S(O)₂Me,
-CH₂S(O)₂Et,
-CH₂S(O)iPr,
-CH₂S(O)₂iPr,
-CH₂S(O)tBu,
-CH₂S(O)₂tBu,
-CH₂CO₂H, CH₂C(O)NH₂,
-CH₂C(O)NMe₂,
-CH₂C(O)NHMe,
-CH₂C(O)-N-pyrrolidine,
-CH₂S(O)₂Me, CH₂S(O)Me,
-CH(OH)CO₂H,
-CH(OH)C(O)NH₂,
-CH(OH)C(O)NHMe,
-CH(OH)C(O)NMe₂,
-CH(OH)C(O)NEt₂,
-CH₂CH₂CO₂H,
-CH₂CH₂CO₂Me,
-CH₂CH₂CO₂Et,
-CH₂CH₂C(O)NH₂,
-CH₂CH₂C(O)NHMe,
-CH₂CH₂C(O)NMe₂,
-CH₂CH₂-5-tetrazolyl,

-CH₂CH₂S(O)₂Me,
-CH₂CH₂S(O)Me,
-CH₂CH₂S(O)₂Et,
-CH₂CH₂S(O) Et,
-CH₂CH₂S(O)iPr,
-CH₂CH₂S(O)₂iPr,
-CH₂CH₂S(O)tBu,
-CH₂CH₂S(O)₂tBu,
-CH₂CH₂S(O)NH₂,
-CH₂CH₂S(O)NHMe,
-CH₂CH₂S(O)NMe₂,
-CH₂CH₂S(O)₂NH₂,
-CH₂CH₂S(O)₂NHMe
-CH₂CH₂S(O)₂NMe₂,
-CH₂CH₂CH₂S(O)Me,
-CH₂CH₂CH₂S(O)Et,
-CH₂CH₂CH₂S(O)₂Me,
-CH₂CH₂CH₂S(O)₂Et,
-C(O)OH,
-5-tetrazolyl,






provided that RB is substituted at either the 6 or 7 position of the benzothiophene ring, except that RB is substituted only at the 7 position of the benzothiophene ring when the group $-(L_{TB})-Z_{TB}$ is at the 6 position.; and

provided that RB' is substituted at either the 4 or 5 position of the benzothiophene ring, except that RB' is substituted only at the 5 position of the benzothiophene ring when the group $-(L_{TB})-Z_{TB}$ is at the 6 position of the phenyl ring; and

provided that RP is substituted at either the 2, or 5 or 6 position of the phenyl ring.

2. . A compound or a pharmaceutically acceptable salt or a prodrug derivative thereof represented by formula (IB):



wherein





R and R' are independently C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, or together R and R' form a substituted or unsubstituted, saturated or unsaturated carbocyclic ring having from 3 to 8 carbon atoms;

RP, RB₄, RT₃, and RB are independently selected from the group consisting of hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, -O-C₁-C₅ alkyl, -S-C₁-C₅ alkyl, -O-C₁-C₅ fluoroalkyl, -CN, -NO₂, acetyl, -S-C₁-C₅ fluoroalkyl, C₂-C₅ alkenyl, C₃-C₅ cycloalkyl, and C₃-C₅ cycloalkenyl;

RP₃ and RB₇ are independently selected from hydrogen, halo, C₁-C₅ alkyl, C₁-C₅ fluoroalkyl, -O-C₁-C₅ alkyl, -S-C₁-C₅ alkyl, -O-C₁-C₅ fluoroalkyl, -CN, -NO₂, acetyl, -S-C₁-C₅ fluoroalkyl, C₂-C₅ alkenyl, C₃-C₅ cycloalkyl, or C₃-C₅ cycloalkenyl;


(L_{P1}), (L_{P2}), and (L_{BT}) are divalent linking groups independently selected from the group consisting of

a bond






R40

R40



O



where m is 0, 1, or 2, and each R40 is independently hydrogen, C₁-C₅ alkyl, or C₁-C₅ fluoroalkyl;

Z_P is

branched C₃-C₅ alkyl,

3-methyl-3-hydroxypentyl,

3-methyl-3-hydroxypentenyl,

3-methyl-3-hydroxypentynyl,

3-ethyl-3-hydroxypentyl,

3-ethyl-3-hydroxypentenyl,

3-ethyl-3-hydroxypentynyl,
3-ethyl-3-hydroxy-4-methylpentyl,
3-ethyl-3-hydroxy-4-methylpentenyl,
3-ethyl-3-hydroxy-4-methylpentynyl,
3-propyl-3-hydroxypentyl,
3-propyl-3-hydroxypentenyl,
3-propyl-3-hydroxypentynyl,
1-hydroxy-2-methyl-1-(methylethyl)propyl,
3-methyl-3-hydroxy-4,4-dimethylpentyl,
3-methyl-3-hydroxy-4,4-dimethylpentenyl,
3-methyl-3-hydroxy-4,4-dimethylpentyl,
3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
3-ethyl-3-hydroxy-4,4-dimethylpentenyl,
3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
1-hydroxycyclopentenyl,
1-hydroxycyclohexenyl,
1-hydroxycycloheptenyl,
1-hydroxycyclooctenyl,
1-hydroxycyclopropyl,
1-hydroxycyclobutyl,
1-hydroxycyclopentyl,
1-hydroxycyclohexyl,
1-hydroxycycloheptyl, or
1-hydroxycyclooctyl;

provided, however, that when

Z_P is

3-methyl-3-hydroxypentyl,
3-methyl-3-hydroxypentenyl,
3-methyl-3-hydroxypentynyl,
3-ethyl-3-hydroxypentyl,
3-ethyl-3-hydroxypentenyl,
3-ethyl-3-hydroxypentynyl,
3-ethyl-3-hydroxy-4-methylpentyl,
3-ethyl-3-hydroxy-4-methylpentenyl,

3-ethyl-3-hydroxy-4-methylpentynyl,
 3-propyl-3-hydroxypentyl,
 3-propyl-3-hydroxypentenyl,
 3-propyl-3-hydroxypentynyl,
 3-methyl-3-hydroxy-4,4-dimethylpentyl,
 3-methyl-3-hydroxy-4,4-dimethylpentenyl,
 3-methyl-3-hydroxy-4,4-dimethylpentyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentynyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentenyl,
 3-ethyl-3-hydroxy-4,4-dimethylpentynyl, or
 1-hydroxy-2-methyl-1-(methylethyl)propyl;

then (L_{P1}) and (L_{P2}) combine as a bond;

Z_{BT} is selected from

-O-(C₁-C₅ alkyl),
 -O-(C₂-C₅ alkenyl),
 -O-(C₃-C₅ cycloalkyl),
 -O-(C₃-C₅ cycloalkenyl),
 -O-(C₁-C₅ hydroxyalkyl),
 -O-(C₁-C₅ fluoroalkyl),
 -O-(C₁-C₅ alkyl)-phenyl,
 -O-(C₁-C₅ alkyl)-(O)-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl) NH₂,
 -O-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl)₂,
 -O-(C₁-C₅ alkyl)-C(O)-NH₂,
 -O-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
 -O-(C₁-C₅ alkyl)-C(O)-OH,
 -O-(C₁-C₅ alkyl)-C(O)-NH-5-tetrazolyl,
 -O-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
 -O-(C₁-C₅ alkyl)-NH₂,

-O-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-O-(C₁-C₅ alkyl)-N-pyrrolidine,
-O-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-O-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-SO₂-NH₂,
-O-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-NH₂,
-O-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-O-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-O-(C₁-C₅ alkyl)-5-tetrazolyl,
-O-CH₂-CO₂H,
-O-CH₂-5-tetrazolyl,
-O-(C₁-C₅ alkyl),
-O-C(O)-NH₂,
-O-C(O)-N-(CH₃)₂,
-O-C(S)-N-(CH₃)₂,
-O-C(O)-O-(C₁-C₅ alkyl),
-O-(5-tetrazolyl),
-O-SO₂-(C₁-C₅ alkyl),
-O-SO₂-NH₂,
-O-SO₂-NH-(C₁-C₅ alkyl),
-O-SO₂-N-(C₁-C₅ alkyl)₂,

-O-S(O)-(C₁-C₅ alkyl),
-O-S(O)-NH₂,
-O-S(O)-NH-(C₁-C₅ alkyl),
-O-S(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl),
-S-(C₂-C₅ alkenyl),
-S-(C₃-C₅ cycloalkyl),
-S-(C₃-C₅ cycloalkenyl),
-S-(C₁-C₅ fluoroalkyl),
-S-(C₁-C₅ hydroxyalkyl),
-S-(C₁-C₅ alkyl)-phenyl,
-S-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-OH,
-S-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-NH₂,
-S-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl) NH₂,
-S-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-S-(C₁-C₅ alkyl)-N-pyrrolidine,
-S-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-S-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-SO₂-NH₂,
-S-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,

-S-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-5-tetrazolyl,
-S-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-S(O)-NH₂,
-S-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-S-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-S-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₂-C₅ alkenyl),
-SO₂-(C₃-C₅ cycloalkyl),
-SO₂-(C₃-C₅ cycloalkenyl),
-SO₂-(C₁-C₅ hydroxyalkyl),
-SO₂-(C₁-C₅ fluoroalkyl),
-SO₂-(C₁-C₅)-phenyl,
-SO₂-NH₂,
-SO₂-NH-(C₁-C₅ alkyl),
-SO₂-NH-CH₂-C(O)OH,
-SO₂-NH-CH₂-C(O)(O-C₁-C₅ alkyl),
-SO₂-NH-(C₁-C₅ alkyl)-C(O)OH,
-SO₂-NH-(C₁-C₅ alkyl)-C(O)(O-C₁-C₅ alkyl),
-SO₂-NHC(O)-(C₃-C₆ cycloalkyl),
-SO₂-NH-C(O)-(C₁-C₅ alkyl),
-SO₂-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl) NH₂,
-SO₂-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,

-SO₂-(C₁-C₅ alkyl)-C(O)-NH₂,
-SO₂-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-SO₂-(C₁-C₅ alkyl)-N-pyrrolidine,
-SO₂-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-SO₂-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-C(O)-OH,
-SO₂-(C₁-C₅ alkyl)-5-tetrazolyl,
-SO₂-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-SO₂-NH₂,
-SO₂-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-SO₂-(C₁-C₅ alkyl),
-SO₂-(C₂-C₅ alkenyl),
-SO₂-(C₃-C₅ cycloalkyl),
-SO₂-(C₃-C₅ cycloalkenyl),
-SO₂-(C₁-C₅ hydroxyalkyl),
-SO₂-(C₁-C₅ fluoroalkyl),
-SO₂-(C₁-C₅)-phenyl,
-SO₂-N=CHN(C₁-C₅ alkyl)₂,
-S(O)-NH₂,
-S(O)-NH-(C₁-C₅ alkyl),
-S(O)-NH-CH₂-C(O)OH
-S(O)-NH-(C₁-C₅ alkyl)-C(O)OH,
-S(O)-NH-CH₂-C(O)(O-C₁-C₅ alkyl),

-S(O)-NH-(C₁-C₅ alkyl)-C(O)(O-C₁-C₅ alkyl),
-S(O)HC(O)-(C₃-C₆ cycloalkyl),
-S(O)-NH-C(O)-(C₁-C₅ alkyl),
-S(O)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-C(O)-NH₂,
-S(O)-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-NH-S(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-S(O)-(C₁-C₅ alkyl)-N-pyrrolidine,
-S(O)-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-3-yl),
-S(O)-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-C(O)-OH,
-S(O)-(C₁-C₅ alkyl)-5-tetrazolyl,
-S(O)-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-SO₂-NH₂,
-S(O)-(C₁-C₅ alkyl)-S(O)-NH₂,
-S(O)-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-S(O)-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-S(O)-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),

-S(O)-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
 -S(O)-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
 -S(O)-N=CHN(C₁-C₅ alkyl)₂,
 -NHC(S)NH₂,
 -NHC(S)NH-(C₁-C₅ alkyl),
 -NHC(S)N-(C₁-C₅ alkyl)₂,
 -NHC(S)NH-(C₂-C₅ alkenyl),
 -NHC(S)NH-(C₃-C₅ cycloalkyl),
 -NHC(S)NH-(C₃-C₅ cycloalkenyl),
 -NHC(S)NH-(C₁-C₅ fluoroalkyl),
 -NHC(S)NH-C₁-C₅ hydroxyalkyl,
 -NHC(S)NH-(C₁-C₅ fluoroalkyl)
 -NHC(S)NH-phenyl,
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-OH,
 -NHC(S)NH-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-(O-C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-NH₂,
 -NHC(S)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-NH₂,
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,
 -NHC(S)NH-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-NH-S(O)-(C₁-C₅ alkyl),
 -NHC(S)NH-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
 -NHC(S)NH-(C₁-C₅ alkyl)-N-pyrrolidine,
 -NHC(S)NH-(C₁-C₅ alkyl)-(1-methylpyrrolidin-2-one-
 3-yl),

-NHC(S)NH-(C₁-C₅ alkyl)-5-tetrazolyl,
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-NH₂,
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-NH-(C₁-C₅ alkyl),
-NHC(S)NH-(C₁-C₅ alkyl)-S(O)-N-(C₁-C₅ alkyl)₂,
-NHC(S)NH-(C₁-C₅ alkyl)-P(O)-(O-C₁-C₅ alkyl)₂,
-NHC(O)NH₂,
-NHC(O)NH-(C₁-C₅ alkyl),
-NHC(O)N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₂-C₅ alkenyl),
-NHC(O)NH-(C₃-C₅ cycloalkyl),
-NHC(O)NH-(C₃-C₅ cycloalkenyl),
-NHC(O)NH-(C₁-C₅ hydroxyalkyl),
-NHC(O)NH-(C₁-C₅ fluoroalkyl),
-NHC(O)NH-phenyl,
-NHC(O)NH-(C₁-C₅ alkyl)-NH₂,
-NHC(O)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₁-C₅ alkyl)-O-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-NH₂,
-NHC(O)NH-(C₁-C₅ alkyl)-NH-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-NH₂,
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-NH-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-N-(C₁-C₅ alkyl)₂,

-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-NH-SO₂-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-N-pyrrolidin-2-one,
-NHC(O)NH-(C₁-C₅ alkyl)-N-pyrrolidine,
-NHC(O)NH-(C₁-C₅ alkyl)-
 (1-methylpyrrolidin-2-one-3-yl),
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-OH,
-NHC(O)NH-(C₁-C₅ alkyl)-C(O)-O-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-5-tetrazolyl,
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-NH₂,
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-NH-(C₁-C₅ alkyl),
-NHC(O)NH-(C₁-C₅ alkyl)-SO₂-N-(C₁-C₅ alkyl)₂,
-NHC(O)NH-(C₁-C₅ alkyl)-P(O)-O-(C₁-C₅ alkyl)₂ ,
-NH₂,
-NH-(C₁-C₅ alkyl),
-NH-CH₂-C(O)OH,
-N-(C₁-C₅ alkyl)₂,
-NH-C(O)-NH₂,
-NH-C(O)-NH-(C₁-C₅ alkyl),
-NH-C(O)-N-(C₁-C₅ alkyl)₂,
-NH-C(O)-(C₁-C₅ alkyl),
-NH-SO₂-(C₁-C₅ alkyl),
-NH-S(O)-(C₁-C₅ alkyl),
-N(CH₃)(OCH₃),
-N(OH)(CH₃),
-N-pyrrolidin-2-one,
-N-pyrrolidine,
-(1-methylpyrrolidin-2-one-3-yl),

-CO₂H,
-CO₂Me,
-CO₂Et,
-C(O)CH₂S(O)Me,
-C(O)CH₂S(O)Et,
-C(O)CH₂S(O)₂Me,
-C(O)CH₂S(O)₂Et,
-C(O)CH₂CH₂S(O)Me,
-C(O)CH₂CH₂S(O)Et,
-C(O)CH₂CH₂S(O)₂Me,
-C(O)CH₂CH₂S(O)₂Et,
-C(O)CH(Me)CH₂CO₂H,
-C(O)CH(Me)CH₂CO₂Me,
-C(O)CH(Me)CH₂CO₂Et,
-C(O)CH(Me)CH₂CO₂iPr,
-C(O)CH(Me)CH₂CO₂tBu,
-C(O)CH(Me)CH(Me)CO₂H,
-C(O)CH(Me)CH(Me)CO₂Me,
-C(O)CH(Me)CH(Me)CO₂Et,
-C(O)CH(Me)CH(Me)CO₂iPr,
-C(O)CH(Me)CH(Me)CO₂tBu,
-C(O)CH(Me)C(Me)₂CO₂H,
-C(O)CH(Me)C(Me)₂CO₂Me,
-C(O)CH(Me)C(Me)₂CO₂Et,
-C(O)CH(Me)C(Me)₂CO₂iPr,
-C(O)CH(Me)C(Me)₂CO₂tBu,
-C(O)CH(Me)CH(Et)CO₂H,
-C(O)CH(Me)CH(Et)CO₂Me,
-C(O)CH(Me)CH(Et)CO₂Et,










-C(O)CH(Me)CH(Et)CO₂iPr,
-C(O)CH(Me)CH(Et)CO₂tBu,
-C(O)C(O)OH,
-C(O)C(O)NH₂,
-C(O)C(O)NHMe,
-C(O)C(O)NMe₂,
-C(O)NH₂,
-C(O)NMe₂,
-C(O)NH-CH₂-C(O)OH,
-C(O)NH-CH₂-C(O)OMe,
-C(O)NH-CH₂-C(O)OEt,
-C(O)NH-CH₂-C(O)OiPr,
-C(O)NH-CH₂-C(O)OtBu,
-C(O)NH-CH(Me)-C(O)OH,
-C(O)NH-CH(Me)-C(O)OMe,
-C(O)NH-CH(Me)-C(O)OEt,
-C(O)NH-CH(Me)-C(O)iPr,
-C(O)NH-CH(Me)-C(O)tBu,
-C(O)NH-CH(Et)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OMe,
-C(O)NH-C(Me)₂-C(O)OEt,
-C(O)NH-C(Me)₂-C(O)iPr,
-C(O)NH-C(Me)₂-C(O)tBu,
-C(O)NH-CMe(Et)-C(O)OH,
-C(O)NH-CH(F)-C(O)OH,
-C(O)NH-CH(CF₃)-C(O)OH,
-C(O)NH-CH(OH)-C(O)OH,
-C(O)NH-CH(cyclopropyl)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-CF(Me)-C(O)OH,
-C(O)NH-C(Me)(CF₃)-C(O)OH,
















-C(O)NH-C(Me)(OH)-C(O)OH,
-C(O)NH-C(Me)(cyclopropyl)CO₂H,
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH₂-C(O)OMe,
-C(O)NMe-CH₂-C(O)OEt,
-C(O)NMe-CH₂-C(O)OiPr,
-C(O)NMe-CH₂-C(O)tBu,
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH(Me)-C(O)OH,
-C(O)NMe-CH(F)-C(O)OH,
-C(O)NMe-CH(CF₃)-C(O)OH,
-C(O)NMe-CH(OH)-C(O)OH,
-C(O)NMe-CH(cyclopropyl)-C(O)OH,
-C(O)NMe-C(Me)₂-C(O)OH,
-C(O)NMe-CF(Me)-C(O)OH,
-C(O)NMe-C(Me)(CF₃)-C(O)OH,
-C(O)NMe-C(Me)(OH)-C(O)OH,
-C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,
-C(O)NHS(O)Me,
-C(O)NHSO₂Me,
-C(O)-NH-5-tetrazolyl,
-C(O)NHS(O)Me,
-C(O)NHS(O)Et,
-C(O)NHSO₂Me,
-C(O)NHSO₂Et,
-C(O)NHS(O)iPr,
-C(O)NHSO₂iPr,
-C(O)NHS(O)tBu,
-C(O)NHSO₂tBu,
-C(O)NHCH₂S(O)Me,
-C(O)NHCH₂S(O)Et,
-C(O)NHCH₂SO₂Me,

-C(O)NHCH₂SO₂Et,
-C(O)NHCH₂CH₂S(O)Me,
-C(O)NHCH₂CH₂S(O)Et,
-C(O)NHCH₂CH₂SO₂Me,
-C(O)NHCH₂CH₂SO₂Et,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)SO₂Me,
-C(O)-N(Me)-5-tetrazolyl,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)S(O)Et,
-C(O)N(Me)SO₂Me,
-C(O)N(Me)SO₂Et,
-C(O)N(Me)S(O)iPr,
-C(O)N(Me))SO₂iPr,
-C(O)N(Me))S(O)tBu,
-C(O)N(Me)SO₂tBu,
-C(O)N(Me)CH₂S(O)Me,
-C(O)N(Me)CH₂S(O)Et,
-C(O)N(Me)CH₂SO₂Me,
-C(O)N(Me)CH₂SO₂Et,
-C(O)N(Me)CH₂CH₂S(O)Me,
-C(O)N(Me)CH₂CH₂S(O)Et,
-C(O)N(Me)CH₂CH₂SO₂Me,
-C(O)N(Me)CH₂CH₂SO₂Et,
-CH₂CO₂H,
-CH₂-5-tetrazolyl,
-CH₂CO₂Me,

-CH₂CO₂Et,
-CH₂NHS(O)Me,
-CH₂NHS(O)Et,
-CH₂NHSO₂Me,
-CH₂NHSO₂Et,
-CH₂NHS(O)iPr,
-CH₂NHSO₂iPr,
-CH₂NHS(O)tBu,
-CH₂NHSO₂tBu,
-CH₂NHCH₂CH₂SO₂CH₃,
-CH₂NH(CH₂CO₂H),
-CH₂N(C(O)Me)(CH₂CO₂H),
-CH₂-N-pyrrolidin-2-one,
-CH₂-(1-methylpyrrolidin-2-one-3-yl),
-CH₂S(O)Me,
-CH₂S(O)Et,
-CH₂S(O)₂Me,
-CH₂S(O)₂Et,
-CH₂S(O)iPr,
-CH₂S(O)₂iPr,
-CH₂S(O)tBu,
-CH₂S(O)₂tBu,
-CH₂CO₂H, CH₂C(O)NH₂,
-CH₂C(O)NMe₂,
-CH₂C(O)NHMe,
-CH₂C(O)-N-pyrrolidine,
-CH₂S(O)₂Me, CH₂S(O)Me,
-CH(OH)CO₂H,
-CH(OH)C(O)NH₂,

-CH(OH)C(O)NHMe,
-CH(OH)C(O)NMe₂,
-CH(OH)C(O)NEt₂,
-CH₂CH₂CO₂H,
-CH₂CH₂CO₂Me,
-CH₂CH₂CO₂Et,
-CH₂CH₂C(O)NH₂,
-CH₂CH₂C(O)NHMe,
-CH₂CH₂C(O)NMe₂,
-CH₂CH₂-5-tetrazolyl,
-CH₂CH₂S(O)₂Me,
-CH₂CH₂S(O)Me,
-CH₂CH₂S(O)₂Et,
-CH₂CH₂S(O) Et,
-CH₂CH₂S(O)iPr,
-CH₂CH₂S(O)₂iPr,
-CH₂CH₂S(O)tBu,
-CH₂CH₂S(O)₂tBu,
-CH₂CH₂S(O)NH₂,
-CH₂CH₂S(O)NHMe,
-CH₂CH₂S(O)NMe₂,
-CH₂CH₂S(O)₂NH₂,
-CH₂CH₂S(O)₂NHMe,
-CH₂CH₂S(O)₂NMe₂,
-CH₂CH₂CH₂S(Θ)Me,
-CH₂CH₂CH₂S(O)Et,
-CH₂CH₂CH₂S(O)₂Me,
-CH₂CH₂CH₂S(O)₂Et,
-C(O)OH,
-5-tetrazolyl,





-1,3,4-oxadiazolin-2-one-5-yl,

-imidazolidine-2,4-dione-5-yl,
-isoxazol-3-ol-yl, or
-1,3,4-oxadiazolin-2-thione-5-yl;

provided that RP is substituted at either the 2, 5, or 6 position of the phenyl ring.

3. The compound of Claim 1

wherein for Formula IA;

R and R' are independently methyl or ethyl;

RP and RT₃ are independently, hydrogen or methyl;

RP₃ and RB are independently hydrogen, methyl, ethyl, -O-methyl, or cyclopropyl;

(L_{P1}) and (L_{TB}) divalent linking groups are both bonds;

(L_{P2}) is a bond, -CH₂-, -CH(OH)-, or -C(Me)OH-;

Z_P is 1,1-dimethylethyl; 1-hydroxycyclopentyl, 1-hydroxycyclohexyl, 3-ethyl-3-hydroxypentyl, 3-ethyl-3-hydroxypentenyl, 3-ethyl-3-hydroxypentynyl;

Z_{TB} is

-CO₂H,
-CO₂Me,
-CO₂Et,
-C(O)CH₂S(O)Me,
-C(O)CH₂S(O)Et,
-C(O)CH₂S(O)₂Me,
-C(O)CH₂S(O)₂Et,
-C(O)CH₂CH₂S(O)Me,
-C(O)CH₂CH₂S(O)Et,
-C(O)CH₂CH₂S(O)₂Me,
-C(O)CH₂CH₂S(O)₂Et,
-C(O)CH(Me)CH₂CO₂H,
-C(O)CH(Me)CH₂CO₂Me,

-C(O)CH(Me)CH₂CO₂Et,
-C(O)CH(Me)CH₂CO₂iPr,
-C(O)CH(Me)CH₂CO₂tBu,
-C(O)CH(Me)CH(Me)CO₂H,
-C(O)CH(Me)CH(Me)CO₂Me,
-C(O)CH(Me)CH(Me)CO₂Et,
-C(O)CH(Me)CH(Me)CO₂iPr,
-C(O)CH(Me)CH(Me)CO₂tBu,
-C(O)CH(Me)C(Me)₂CO₂H,
-C(O)CH(Me)C(Me)₂CO₂Me,
-C(O)CH(Me)C(Me)₂CO₂Et,
-C(O)CH(Me)C(Me)₂CO₂iPr,
-C(O)CH(Me)C(Me)₂CO₂tBu,
-C(O)CH(Me)CH(Et)CO₂H,
-C(O)CH(Me)CH(Et)CO₂Me,
-C(O)CH(Me)CH(Et)CO₂Et,
-C(O)CH(Me)CH(Et)CO₂iPr,
-C(O)CH(Me)CH(Et)CO₂tBu,
-C(O)C(O)OH,
-C(O)C(O)NH₂,
-C(O)C(O)NHMe,
-C(O)C(O)NMe₂,
-C(O)NH₂,
-C(O)NMe₂,
-C(O)NH-CH₂-C(O)OH,
-C(O)NH-CH₂-C(O)OMe,
-C(O)NH-CH₂-C(O)OEt,
-C(O)NH-CH₂-C(O)OiPr,
-C(O)NH-CH₂-C(O)OtBu,
-C(O)NH-CH(Me)-C(O)OH,

-C(O)NH-CH(Me)-C(O)OMe,
-C(O)NH-CH(Me)-C(O)OEt,
-C(O)NH-CH(Me)-C(O)iPr,
-C(O)NH-CH(Me)-C(O)tBu,
-C(O)NH-CH(Et)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OMe,
-C(O)NH-C(Me)₂-C(O)OEt,
-C(O)NH-C(Me)₂-C(O)iPr,
-C(O)NH-C(Me)₂-C(O)tBu,
-C(O)NH-CMe(Et)-C(O)OH,
-C(O)NH-CH(F)-C(O)OH,
-C(O)NH-CH(CF₃)-C(O)OH,
-C(O)NH-CH(OH)-C(O)OH,
-C(O)NH-CH(cyclopropyl)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-CF(Me)-C(O)OH,
-C(O)NH-C(Me)(CF₃)-C(O)OH,
-C(O)NH-C(Me)(OH)-C(O)OH,
-C(O)NH-C(Me)(cyclopropyl)CO₂H
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH₂-C(O)OMe,
-C(O)NMe-CH₂-C(O)OEt,
-C(O)NMe-CH₂-C(O)OiPr,
-C(O)NMe-CH₂-C(O)tBu,
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH(Me)-C(O)OH,
-C(O)NMe-CH(F)-C(O)OH,
-C(O)NMe-CH(CF₃)-C(O)OH,
-C(O)NMe-CH(OH)-C(O)OH,
-C(O)NMe-CH(cyclopropyl)-C(O)OH,
-C(O)NMe-C(Me)₂-C(O)OH,

-C(O)NMe-CF(Me)-C(O)OH,
-C(O)NMe-C(Me)(CF₃)-C(O)OH,
-C(O)NMe-C(Me)(OH)-C(O)OH,
-C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,
-C(O)NHS(O)Me,
-C(O)NHSO₂Me,
-C(O)-NH-5-tetrazolyl,
-C(O)NHS(O)Me,
-C(O)NHS(O)Et,
-C(O)NHSO₂Me,
-C(O)NHSO₂Et,
-C(O)NHS(O)iPr,
-C(O)NHSO₂iPr,
-C(O)NHS(O)tBu,
-C(O)NHSO₂tBu,
-C(O)NHCH₂S(O)Me,
-C(O)NHCH₂S(O)Et,
-C(O)NHCH₂SO₂Me,
-C(O)NHCH₂SO₂Et,
-C(O)NHCH₂CH₂S(O)Me,
-C(O)NHCH₂CH₂S(O)Et,
-C(O)NHCH₂CH₂SO₂Me,
-C(O)NHCH₂CH₂SO₂Et,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)SO₂Me,
-C(O)-N(Me)-5-tetrazolyl,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)S(O)Et,
-C(O)N(Me)SO₂Me,
-C(O)N(Me)SO₂Et,
-C(O)N(Me)S(O)iPr,

-C(O)N(Me))SO₂iPr,
-C(O)N(Me))S(O)tBu,
-C(O)N(Me)SO₂tBu,
-C(O)N(Me)CH₂S(O)Me,
-C(O)N(Me)CH₂S(O)Et,
-C(O)N(Me)CH₂SO₂Me,
-C(O)N(Me)CH₂SO₂Et,
-C(O)N(Me)CH₂CH₂S(O)Me,
-C(O)N(Me)CH₂CH₂S(O)Et,
-C(O)N(Me)CH₂CH₂SO₂Me,
-C(O)N(Me)CH₂CH₂SO₂Et,
-CH₂CO₂H,
-CH₂-5-tetrazolyl,
-CH₂CO₂Me,
-CH₂CO₂Et,
-CH₂NHS(O)Me,
-CH₂NHS(O)Et,
-CH₂NHSO₂Me,
-CH₂NHSO₂Et,
-CH₂NHS(O)iPr,
-CH₂NHSO₂iPr,
-CH₂NHS(O)tBu,
-CH₂NHSO₂tBu,
-CH₂NHCH₂CH₂SO₂CH₃,
-CH₂NH(CH₂CO₂H),
-CH₂N(C(O)Me)(CH₂CO₂H),
-CH₂-N-pyrrolidin-2-one,
-CH₂-(1-methylpyrrolidin-2-one-3-yl),
-CH₂S(O)Me,

-CH₂S(O)Et,
-CH₂S(O)₂Me,
-CH₂S(O)₂Et,
-CH₂S(O)iPr,
-CH₂S(O)₂iPr,
-CH₂S(O)tBu,
-CH₂S(O)₂tBu,
-CH₂CO₂H, CH₂C(O)NH₂,
-CH₂C(O)NMe₂,
-CH₂C(O)NHMe,
-CH₂C(O)-N-pyrrolidine,
-CH₂S(O)₂Me, CH₂S(O)Me,
-CH(OH)CO₂H,
-CH(OH)C(O)NH₂,
-CH(OH)C(O)NHMe,
-CH(OH)C(O)NMe₂,
-CH(OH)C(O)NEt₂,
-CH₂CH₂CO₂H,
-CH₂CH₂CO₂Me,
-CH₂CH₂CO₂Et,
-CH₂CH₂C(O)NH₂,
-CH₂CH₂C(O)NHMe,
-CH₂CH₂C(O)NMe₂,
-CH₂CH₂-5-tetrazolyl,
-CH₂CH₂S(O)₂Me,
-CH₂CH₂S(O)Me,
-CH₂CH₂S(O)₂Et,
-CH₂CH₂S(O)Et,
-CH₂CH₂S(O)iPr,

$-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{iPr}$,
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{tBu}$,
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{tBu}$,
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{NH}_2$,
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{NHMe}$,
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{NMe}_2$,
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{NH}_2$,
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{NHMe}$,
 $-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{NMe}_2$,
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{Me}$,
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{Et}$,
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{Me}$, or
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{Et}$.

4. The compound of claim 2 wherein for formula IB;

R and R' are independently methyl or ethyl;

RP, RB, RB₄, and RT₃ are independently, hydrogen or methyl;

RP₃ and RB₇ are independently hydrogen, methyl, ethyl, -O-methyl, or cyclopropyl;

(L_{P1}) and (L_{BT}) divalent linking groups are both bonds;

(L_{P2}) is a bond, -CH₂-, -CH(OH)-, or -C(Me)OH-;

Z_P is 1,1-dimethylethyl; 1-hydroxycyclopentyl, 1-hydroxycyclohexyl, 3-ethyl-3-hydroxypentyl, 3-ethyl-3-hydroxypentenyl, 3-ethyl-3-hydroxypentynyl;

Z_{BT} is

$-\text{CO}_2\text{H}$,
 $-\text{CO}_2\text{Me}$,
 $-\text{CO}_2\text{Et}$,
 $-\text{C}(\text{O})\text{CH}_2\text{S}(\text{O})\text{Me}$,
 $-\text{C}(\text{O})\text{CH}_2\text{S}(\text{O})\text{Et}$,
 $-\text{C}(\text{O})\text{CH}_2\text{S}(\text{O})_2\text{Me}$,
 $-\text{C}(\text{O})\text{CH}_2\text{S}(\text{O})_2\text{Et}$,

-C(O)CH₂CH₂S(O)Me,
-C(O)CH₂CH₂S(O)Et,
-C(O)CH₂CH₂S(O)₂Me,
-C(O)CH₂CH₂S(O)₂Et,
-C(O)CH(Me)CH₂CO₂H,
-C(O)CH(Me)CH₂CO₂Me,
-C(O)CH(Me)CH₂CO₂Et,
-C(O)CH(Me)CH₂CO₂iPr,
-C(O)CH(Me)CH₂CO₂tBu,
-C(O)CH(Me)CH(Me)CO₂H,
-C(O)CH(Me)CH(Me)CO₂Me,
-C(O)CH(Me)CH(Me)CO₂Et,
-C(O)CH(Me)CH(Me)CO₂iPr,
-C(O)CH(Me)CH(Me)CO₂tBu,
-C(O)CH(Me)C(Me)₂CO₂H,
-C(O)CH(Me)C(Me)₂CO₂Me,
-C(O)CH(Me)C(Me)₂CO₂Et,
-C(O)CH(Me)C(Me)₂CO₂iPr,
-C(O)CH(Me)C(Me)₂CO₂tBu,
-C(O)CH(Me)CH(Et)CO₂H,
-C(O)CH(Me)CH(Et)CO₂Me,
-C(O)CH(Me)CH(Et)CO₂Et,
-C(O)CH(Me)CH(Et)CO₂iPr,
-C(O)CH(Me)CH(Et)CO₂tBu,
-C(O)C(O)OH,
-C(O)C(O)NH₂,
-C(O)C(O)NHMe,
-C(O)C(O)NMe₂,
-C(O)NH₂,

-C(O)NMe₂,
-C(O)NH-CH₂-C(O)OH,
-C(O)NH-CH₂-C(O)OMe,
-C(O)NH-CH₂-C(O)OEt,
-C(O)NH-CH₂-C(O)OiPr,
-C(O)NH-CH₂-C(O)OtBu,
-C(O)NH-CH(Me)-C(O)OH,
-C(O)NH-CH(Me)-C(O)OMe,
-C(O)NH-CH(Me)-C(O)OEt,
-C(O)NH-CH(Me)-C(O)iPr,
-C(O)NH-CH(Me)-C(O)tBu,
-C(O)NH-CH(Et)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OMe,
-C(O)NH-C(Me)₂-C(O)OEt,
-C(O)NH-C(Me)₂-C(O)iPr,
-C(O)NH-C(Me)₂-C(O)tBu,
-C(O)NH-CMe(Et)-C(O)OH,
-C(O)NH-CH(F)-C(O)OH,
-C(O)NH-CH(CF₃)-C(O)OH,
-C(O)NH-CH(OH)-C(O)OH,
-C(O)NH-CH(cyclopropyl)-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-C(Me)₂-C(O)OH,
-C(O)NH-CF(Me)-C(O)OH,
-C(O)NH-C(Me)(CF₃)-C(O)OH,
-C(O)NH-C(Me)(OH)-C(O)OH,
-C(O)NH-C(Me)(cyclopropyl)CO₂H
-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH₂-C(O)OMe,
-C(O)NMe-CH₂-C(O)OEt,
-C(O)NMe-CH₂-C(O)OiPr,
-C(O)NMe-CH₂-C(O)tBu,

-C(O)NMe-CH₂-C(O)OH,
-C(O)NMe-CH(Me)-C(O)OH,
-C(O)NMe-CH(F)-C(O)OH,
-C(O)NMe-CH(CF₃)-C(O)OH,
-C(O)NMe-CH(OH)-C(O)OH,
-C(O)NMe-CH(cyclopropyl)-C(O)OH,
-C(O)NMe-C(Me)₂-C(O)OH,
-C(O)NMe-CF(Me)-C(O)OH,
-C(O)NMe-C(Me)(CF₃)-C(O)OH,
-C(O)NMe-C(Me)(OH)-C(O)OH,
-C(O)NMe-C(Me)(cyclopropyl)-C(O)OH,
-C(O)NHS(O)Me,
-C(O)NHSO₂Me,
-C(O)-NH-5-tetrazolyl,
-C(O)NHS(O)Me,
-C(O)NHS(O)Et,
-C(O)NHSO₂Me,
-C(O)NHSO₂Et,
-C(O)NHS(O)iPr,
-C(O)NHSO₂iPr,
-C(O)NHS(O)tBu,
-C(O)NHSO₂tBu,
-C(O)NHCH₂S(O)Me,
-C(O)NHCH₂S(O)Et,
-C(O)NHCH₂SO₂Me,
-C(O)NHCH₂SO₂Et,
-C(O)NHCH₂CH₂S(⊖)Me,
-C(O)NHCH₂CH₂S(O)Et,
-C(O)NHCH₂CH₂SO₂Me,
-C(O)NHCH₂CH₂SO₂Et,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)SO₂Me,


-C(O)-N(Me)-5-tetrazolyl,
-C(O)N(Me)S(O)Me,
-C(O)N(Me)S(O)Et,
-C(O)N(Me)SO₂Me,
-C(O)N(Me)SO₂Et,
-C(O)N(Me)S(O)iPr,
-C(O)N(Me))SO₂iPr,
-C(O)N(Me))S(O)tBu,
-C(O)N(Me)SO₂tBu,
-C(O)N(Me)CH₂S(O)Me,
-C(O)N(Me)CH₂S(O)Et,
-C(O)N(Me)CH₂SO₂Me,
-C(O)N(Me)CH₂SO₂Et,
-C(O)N(Me)CH₂CH₂S(O)Me,
-C(O)N(Me)CH₂CH₂S(O)Et,
-C(O)N(Me)CH₂CH₂SO₂Me,
-C(O)N(Me)CH₂CH₂SO₂Et,
-CH₂CO₂H,
-CH₂-5-tetrazolyl,
-CH₂CO₂Me,
-CH₂CO₂Et,
-CH₂NHS(O)Me,
-CH₂NHS(O)Et,
-CH₂NHSO₂Me,
-CH₂NHSO₂Et,
-CH₂NHS(O)iPr,
-CH₂NHSO₂iPr,
-CH₂NHS(O)tBu,
-CH₂NHSO₂tBu,
-CH₂NHCH₂CH₂SO₂CH₃,

-CH₂NH(CH₂CO₂H),
-CH₂N(C(O)Me)(CH₂CO₂H),
-CH₂-N-pyrrolidin-2-one,
-CH₂-(1-methylpyrrolidin-2-one-3-yl),
-CH₂S(O)Me,
-CH₂S(O)Et,
-CH₂S(O)₂Me,
-CH₂S(O)₂Et,
-CH₂S(O)iPr,
-CH₂S(O)₂iPr,
-CH₂S(O)tBu,
-CH₂S(O)₂tBu,
-CH₂CO₂H, CH₂C(O)NH₂,
-CH₂C(O)NMe₂,
-CH₂C(O)NHMe,
-CH₂C(O)-N-pyrrolidine,
-CH₂S(O)₂Me, CH₂S(O)Me,
-CH(OH)CO₂H,
-CH(OH)C(O)NH₂,
-CH(OH)C(O)NHMe,
-CH(OH)C(O)NMe₂,
-CH(OH)C(O)NEt₂,
-CH₂CH₂CO₂H,
-CH₂CH₂CO₂Me,
-CH₂CH₂CO₂Et,
-CH₂CH₂C(O)NH₂,
-CH₂CH₂C(O)NHMe,
-CH₂CH₂C(O)NMe₂,
-CH₂CH₂-5-tetrazolyl,


-CH₂CH₂S(O)₂Me,
-CH₂CH₂S(O)Me,
-CH₂CH₂S(O)₂Et,
-CH₂CH₂S(O) Et,
-CH₂CH₂S(O)iPr,
-CH₂CH₂S(O)₂iPr,
-CH₂CH₂S(O)tBu,
-CH₂CH₂S(O)₂tBu,
-CH₂CH₂S(O)NH₂,
-CH₂CH₂S(O)NHMe,
-CH₂CH₂S(O)NMe₂,
-CH₂CH₂S(O)₂NH₂,
-CH₂CH₂S(O)₂NHMe
-CH₂CH₂S(O)₂NMe₂,
-CH₂CH₂CH₂S(O)Me,
-CH₂CH₂CH₂S(O)Et,
-CH₂CH₂CH₂S(O)₂Me, or
-CH₂CH₂CH₂S(O)₂Et.

5. The compound represented by formula (C1) to (C22) or a pharmaceutically acceptable salt or prodrug derivative thereof:


C1)




C2)




C3)




C4)




C5)



C6)




C7)




C8)

P-16541


185




C9)




C10)



C11)




C12)




C13)

P-16541


186




C14)




C15)




C16)




C17)




C18)




C19)



C20)




C21)




C22)

188



6 The compound represented by the structural formula AA or a pharmaceutically acceptable salt or prodrug thereof:



7 The prodrug derivative of the compound of claim 1 or 2 in the prodrug is a methyl ester; ethyl ester; N,N-diethylglycolamido ester; or morpholinylethyl ester.

8 The salt derivative of the compound of claim 1 or 2 wherein the salt is sodium or potassium.

9 A pharmaceutical formulation comprising the compound of claim 1 or 2 ether with a pharmaceutically acceptable carrier or diluent.

10 A formulation for treating osteoporosis comprising:

Ingredient (A1): the vitamin D receptor modulator of claim 1 or;

Ingredient (B1):

one or more co-agents selected from the group consisting of:

- a. estrogens,
- b. androgens,
- c. calcium supplements,

- d. vitamin D metabolites,
- e. thiazide diuretics,
- f. calcitonin,
- g. bisphosphonates,
- h. SERMS, and
- i. fluorides; and

Ingredient (C1): optionally, a carrier or diluent.

11 The formulation of claim 10 wherein the weight ratio of (A1) to (B1) is from 10:1 to 1:1000.

12 A formulation for treating psoriasis comprising:

Ingredient (A2): the vitamin D receptor modulator of claim 1 or 2;

Ingredient (B2):

one or more co-agents that are conventional for treatment of psoriasis selected from the group consisting of:

- a. topical glucocorticoids ,
- b. salicylic acid,
- c. crude coal tar; and

Ingredient (C2): optionally, a carrier or diluent.

13. The formulation of claim 12 wherein the weight ratio of (A2) to (B2) is from 1:10 to 1:100000.

14. A method of treating a mammal to prevent or alleviate the pathological effects of Acne, Actinic keratosis, Alopecia , Alzheimer's disease, Bone maintenance in zero gravity, Bone fracture healing, Breast cancer, Chemoprevention of Cancer, Crohn's disease, Colon cancer, Type I diabetes, Host-graft rejection, Hypercalcemia , Type II diabetes, Leukemia, Multiple sclerosis, Myelodysplastic syndrome, Insufficient sebum secretion, Osteomalacia, Osteoporosis, Insufficient dermal firmness, Insufficient dermal hydration, Psoriatic arthritis, Prostate cancer, Psoriasis, Renal osteodystrophy, Rheumatoid arthritis, Scleroderma, Skin cancer, Systemic lupus erythematosus, Skin cell damage from , Mustard vesicants, Ulcerative colitis,

Vitiligo, or Wrinkles; wherein the method comprises administering a pharmaceutically effective amount of at least one compound of claim 1 or 2.

15. The method of claim 14 for the treatment of psoriasis.

16. The method of claim 14 for the treatment of osteoporosis.

17. A method of claim 14 for treating a mammal to prevent or alleviate skin cell damage from Mustard vesicants.

18. A method of treating or preventing disease states mediated by the Vitamin D receptor, wherein a mammal in need thereof is administered a pharmaceutically effective amount of the compound of Claim 1 or 2.

19. A compound as claimed in any one of Claims 1 or 2 for use in treating a mammal to prevent or alleviate the pathological effects of Acne, Actinic keratosis, Alopecia, Alzheimer's disease, Bone maintenance in zero gravity, Bone fracture healing, Breast cancer, Chemoprevention of Cancer, Crohn's disease, Colon cancer, Type I diabetes, Host-graft rejection, Hypercalcemia, Type II diabetes, Leukemia, Multiple sclerosis, Myelodysplastic syndrome, Insufficient sebum secretion, Osteomalacia, Osteoporosis, Insufficient dermal firmness, Insufficient dermal hydration, Psoriatic arthritis, Prostate cancer, Psoriasis, Renal osteodystrophy, Rheumatoid arthritis, Scleroderma, Skin cancer, Systemic lupus erythematosus, Skin cell damage from, Mustard vesicants, Ulcerative colitis, Vitiligo, or Wrinkles.

20. A compound as claimed in any one of Claims 1 or 2 for use in treating or preventing disease states mediated by the Vitamin D receptor.

21. A compound as claimed in Claim 1 or 2 substantially as hereinbefore described with reference to any of the Examples.

22. A process for preparing a compound as claimed in claim 1 or 2 substantially as hereinbefore described with reference to any of the Examples.

23. The use of a compound as claimed in claim 1 or 2 substantially as herein described with reference to any of the Assays and Tables for mediating the Vitamin D receptor.

ABSTRACT

The present invention relates to novel, non-secosteroidal, phenyl-benzothiophene compounds with vitamin D receptor (VDR) modulating activity that are less hypercalcemic than $1\alpha,25$ dihydroxy vitamin D₃. These compounds are useful for treating bone disease and psoriasis.

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/037181

International filing date: 16 November 2004 (16.11.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/523,600
Filing date: 20 November 2003 (20.11.2003)

Date of receipt at the International Bureau: 09 December 2004 (09.12.2004)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.